

PCT

PA 1999 00091

PA 1999 00093

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent

(BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE,

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

INTERNATIONAL AFFECATION FUBLISHED UNDER THE FATEIN COOLERATION TREAT (FCT)					
(51) International Patent	1		(11) International Publication Number: WO 99/48380		
A23K 1/165, A01H 5/00, C12N 9/16		A1	(43) International Publication Date: 30 September 1999 (30.09.99)		
(21) International Applica	tion Number: PCT/DK9	99/001:	54 (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB,		
(22) International Filing Date: 22 March 1999 (22.03.99)			GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK,		
(30) Priority Data:			MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW,		
0407/98	23 March 1998 (23.03.98)	D	ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG,		
PA 1998 00806	19 June 1998 (19.06.98)	D	ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ,		
PA 1998 01176	18 September 1998 (18.09.98	3) D	TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI		

DK

DK

(71) Applicant: NOVO NORDISK A/S [DK/DK]; Corporate Patents, Novo Allé, DK-2800 Bagsværd (DK).

22 January 1999 (22.01.99)

22 January 1999 (22.01.99)

(72) Inventor: PETERSEN, Svend; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsværd (DK). SN, TD, TG).

Published

With international search report.

(54) Title: THERMOSTABLE PHYTASES IN FEED PREPARATION AND PLANT EXPRESSION

(57) Abstract

The use of thermostable phytases in the preparation of animal feed, and the expression in plants of such phytases. For preparation of animal feed, a thermostable phytase is added before or during the agglomeration step. Preferred processes are pelleting, extrusion and expansion. A transgenic plant expressing a thermostable phytase may be used directly in animal feed preparation.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

1

Thermostable phytases in feed preparation and plant expression

Technical Field

This application relates to thermostable phytases, viz. their use in processes for the production of animal feed, and their expression in plants.

Background art

WO 91/14782 describes transgenic tobacco and rapeseed plants expressing a phytase derived from Aspergillus ficuum NRRL 3135. The transgenic tobacco seeds are fed to broilers.

US 5,824,779 describes in standard fashion how to produce transgenic alfalfa expressing the same A. ficuum phytase, and the preparation of a phytase-containing concentrate which can be used per se as an animal feed supplement.

EP 0 556 883 Bl describes a method for preparing feed pellets based on an extrusion technique. The addition of temperature sensitive agents, one example of which is phytase, 20 takes place after extrusion of the feed pellets, and the sensitive agents are loaded onto the pellets under reduced pressure.

As acknowledged in EP 0 556 883 Bl the loss of activity of heat-sensitive substances during feed preparation processes is a 25 well-known problem. The above EP-patent proposes to solve this problem by adding these substances under reduced pressure subsequent to the extrusion process. This solution, however, requires a liquid form of the sensitive substance, as well as the installation of additional expensive process equipment.

The present invention provides an improved process for preparing animal feed, as well as improved phytase-expressing transgenic plants.

Summary of the Invention 5

The present invention provides a process of preparing an animal feed, which process comprises an agglomeration of feed ingredients, wherein a thermostable phytase is added before or during the agglomeration.

Also provided is a transgenic plant or part thereof which comprises a DNA-construct encoding a thermostable phytase. 10

The transgenic plant or part thereof, e.g. seeds or leaves, may be used in the feed preparation process of the invention, to thereby provide - in a preferred embodiment - at 15 the same time a nutrient (feed ingredient) and the feed additive phytase.

Brief description of the Figures

In the detailed description of the invention below, 20 reference is made to the drawings, of which

- is a differential scanning calorimetry (DSC) chart Fig. 1 of consensus phytase-1 and consensus phytase-10;
- phytase-10-thermo-Q50T and consensus of DSC Fig. 2 consensus phytase-10-thermo-Q50T-K91A;
- a DSC of consensus phytase-1-thermo[8]-Q50T and 25 Fig. 3 consensus phytase-1-thermo[8]-Q50T-K91A;
 - a DSC of the phytase from A. fumigatus ATCC 13073 Fig. 4 and of its α -mutant; and
- shows the design of the consensus-phytase-1 amino Fig. 5 acid sequence; 30

- Fig. 6 an alignment and the basidiomycete consensus sequence of five Basidiomycete phytases;
- Fig. 7 the design of the consensus-phytase-10 amino acid sequence;
- an alignment for the design of consensus-phytase-11 (all Basidiomycete phytases were used as independent sequences using an assigned vote weight of 0.2 for each Basidiomycete sequence; still further the amino acid sequence of A. niger T213 was used);
- 10 Fig. 9 the DNA and amino acid sequence of consensus-phytase-1-thermo(8)-Q50T-K91A;
 - Fig. 10 the DNA and amino acid sequence of Consensus-phytase-10-thermo(3)-Q50T-K91A;
- Fig. 11 the DNA and amino acid sequence of A. fumigatus ATCC 13073 α -mutant; and
- Fig. 12 the DNA and amino acid sequence of Consensus-phytase-7 which comprises the following mutations as compared to Consensus-phytase-1: S89D, S92G, A94K, D164S, P201S, G203A, G205S, H212P, G224A, D226T, E255T, D256E, V258T, P265S, Q292H, G300K, Y305H, A314T, S364G, M365I, A397S, S398A, G404A, and A405S.

Detailed description of the invention

In the present context a "feed" or an "animal feed" means any natural or artificial diet, meal or the like intended or suitable for being eaten, taken in, digested, by an animal. Food for human beings is included in the above definition of feed.

"Animals" include all animals, be it polygastric animals 30 (ruminants); or monogastric animals such as human beings,

PCT/DK99/0054 WO 99/48380

poultry, swine and fish. Preferred animals are the mono-gastric animals, in particular pigs and broilers.

The concept of "feed ingredients" includes materials from which a feed is to be, or is, produced; or the 5 intended, or actual, component parts of a feed. Feed ingredients for non-human animals are usually, and preferably, selected from amongst the following non-exclusive list:

plant derived products

such as seeds, grains, leaves, roots, tubers, flowers, pods, husks - and they may take the form of flakes, cakes, 10 grits, flour, and the like;

animal derived products

such as fish meal, milk powder, bone extract, meat extract, blood extract and the like;

15 additives

such as minerals, vitamins, aroma compounds, and feed enhancing enzymes.

Phytic acid or myo-inositol 1,2,3,4,5,6-hexakis dihydrogen phosphate (or for short myo-inositol hexakisphosphate) is the 20 primary source of inositol and the primary storage form of phosphate in plant seeds and grains. In the seeds of legumes it accounts for about 70% of the phosphate content. Seeds, cereal grains and legumes are important feed ingredients.

Phytic acid, or its salts phytates - said terms being, otherwise indicated, in the present context used 25 unless synonymously or at random - is an anti-nutritional factor. This is partly due to its binding of nutritionally essential ions such as calcium, trace minerals such as mangane, and also proteins (by electrostatic interaction). And partly due to the 30 fact that the phosphorous thereof is not nutritionally available

either, since phytic acid and its salts, phytates, are often not metabolized.

5

This leads to a need of supplementing food and feed preparations with e.g. inorganic phosphate.

- The non-metabolizable phytic acid phosphorous passes through the gastrointestinal tract of such animals and is excreted with the manure, resulting in an undesirable phosphate pollution of the environment resulting e.g. in eutrophication of the water environment and extensive growth of algae.
- Phytic acid is degradable by phytases. In the present context a "phytase" is an polypeptide or enzyme which exhibits phytase activity, viz. which catalyzes the hydrolysis of phytate (myo-inositol hexakisphosphate) to (1) myo-inositol and/or (2) mono-, di-, tri-, tetra- and/or penta-phosphates thereof and (3) inorganic phosphate.

The production of phytases by plants as well as by microorganisms has been reported. Amongst the microorganisms, phytase producing bacteria as well as phytase producing fungiare known.

There are several descriptions of phytase producing filamentous fungi belonging to the fungal phylum of Ascomycota (ascomycetes). In particular, there are several references to phytase producing ascomycetes of the Aspergillus genus such as Aspergillus terreus (Yamada et al., 1986, Agric. Biol. Chem. 322:1275-1282). Also, the cloning and expression of the phytase gene from Aspergillus niger var. awamori has been described (Piddington et al., 1993, Gene 133:55-62). EP 0420358 describes the cloning and expression of a phytase of Aspergillus ficuum (niger). EP 0684313 describes the cloning and expression of phytases of the ascomycetes Aspergillus niger, Myceliophthora thermophila, Aspergillus terreus. Still further, some partial

6

sequences of phytases of Aspergillus nidulans, Talaromyces thermophilus, Aspergillus fumigatus and another strain of Aspergillus terreus are given.

The cloning and expression of a phytase of Thermomyces 5 lanuginosus is described in WO 97/35017.

WO 98/28409 describes the cloning and expression of several basidiomycete phytases, e.g. from Peniophora lycii, Agrocybe pediades, Paxillus involutus and Trametes pubescens.

According to the Enzyme nomenclature database ExPASy (a 10 repository of information relative to the nomenclature of the the recommendations enzymes primarily based on of Union International Committee of the Nomenclature Biochemistry and Molecular Biology (IUBMB) describing each type of characterized enzyme for which an EC (Enzyme Commission) 15 number has been provided), two different types of phytases are (myo-inositol 3-phytase so-called presently known: Α hexaphosphate 3-phosphohydrolase, EC 3.1.3.8) and a so-called 6phytase (myo-inositol hexaphosphate 6-phosphohydrolase, EC 3.1.3.26). The 3-phytase hydrolyses first the ester bond at a 3-20 position, whereas the 6-phytase hydrolyzes first an ester bond at the 6-position of phytic acid. Both of these types of phytases are included in the above definition of phytase.

Many assays of phytase activity are known, and any of these can be used for the purpose of the present invention. 25 Preferred phytase assays are included in the examples.

The concept of "agglomeration" is defined as a process in which various components are mixed under the influence of heat. The resulting product is preferably an "agglomerate" or conglomerate in which the components adhere to each other while forming a product of a satisfactory physical stability. The formation of dust from such agglomerate is an indication of its

physical stability - the less dust being formed, the better. A suitable assay for dust formation from agglomerates is ASAE standard S 269-1. A satisfactory agglomerate has below 20%, preferably below 15%, more preferably below 10%, even more 5 preferably below 6% dust.

"Under the influence of heat" means that the temperature is at least 65°C, as measured on the product at the outlet of the agglomeration unit. More preferred temperatures are at least 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, or even at least 130°C.

A preferred agglomeration process is operated at an increased pressure. The pressure is typically due to a compacting of the ingredients, optionally in combination with a reduction of the cross-sectional or throughput area. Preferably, by properly adjusting process parameters such as temperature and pressure, the resulting shear forces and shear velocities are of such magnitude, that the starch- and protein-containing feed ingredients become fluid.

"Increased pressure" means increased as compared to normal 20 atmospheric pressure, and the maximum pressure as measured within the agglomeration unit.

The addition of water vapour or steam is often included in agglomeration, but not as an absolute requirement.

Agglomeration includes, but is not limited to, the well25 known processes called extrusion, expansion (or pressure
conditioning) and pelleting (or pellet pressing).

Extrusion is i.a. described at pp. 149-153 of a handbook which is available on request from the Danish Company Sprout-Matador, Glentevej 5-7, DK-6705 Esbjerg Ø or Niels Finsensvej 4, 30 DK-7100 Vejle ("Håndbog i Pilleteringsteknik 1996"). However, in the agglomeration process of the invention, the following

process steps mentioned in the above handbook are entirely optional:

- (i) pre-treating the feed ingredients in a cascade mixer;
- (ii) cutting the product leaving the nozzle-section into pieces
- 5 (iii) of a desired size;
 - (iv) acclimatizing or conditioning it;
 - (v) coating it;
 - (vi) drying it;
 - (vii) cooling it.
- The process of expansion (pressure conditioning) is i.a. described in the same handbook at pp. 61-66. Also for expansion, the above process steps (i)-(vi), in particular steps (i) and (vi), are entirely optional steps.

This is so also for the following process steps:

- 15 (ii') comminuting the product (using e.g. a blade granulator as shown at p. 65);

The process of pelleting is i.a. described in the same 20 handbook at pp. 71-107. Also here, steps (i)-(vii) above are entirely optional steps. These steps are i.a. described in more detail at pp. 29-70 of the above handbook.

In a preferred agglomeration process of the invention, one or more of the above mentioned further process steps (i)-(vii) are included.

A particularly preferred further step is step (i).

In a most preferred embodiment, the feed-ingredients are pre-heated in a first step (a) to a temperature of at least 45°C, preferably at least 50, 55, 60, 65, 70, 75, 80°C; and then heated in a second step (b) to a temperature of at least

65°C, preferably 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, or even at least 130°C.

The addition of thermostable phytase takes place before or during step (a) and/or before or during step (b).

Water is preferably added in step (a). More preferably, heated steam is added during the mixing of the ingredients (steps (a) and/or (b)).

Process step (a) is preferably performed in a cascade mixer (see the above cited handbook p. 44).

A "thermostable" phytase is a phytase which has a Tm (melting temperature) as measured on purified phytase protein by Differential Scanning Calorimetry (DSC) of at least 65°C, preferably using for the DSC a constant heating rate, more preferably of 10°C/min. In preferred embodiments, the Tm is at 15 least 66, 67, 68, 69, 70, 71, 72, 73, 74 or 75°C. Preferably, the Tm is equal to or lower than 150°C, more preferably equal to or lower than 145, 140, 135, 130, 125, 120, 115 or 110°C. Accordingly, preferred intervals of Tm are: 65-150°C, 66-150°C, (etc.) - 75-150°C; 65-145°C, 66-145°C, - (etc.) - 75-145°C; 20 65-140°C, - (etc.) - 75-110°C.

Particularly preferred ranges for Tm are the following: between 65 and 110°C; between 70 and 110°C; between 70 and 100°C; between 75 and 95°C, or between 80 and 90°C.

In Example 3 below, the measurement of Tm by DSC is described, and the Tm's of a number of phytases are shown.

The optimum temperatures are also indicated, since - in the alternative - a thermostable phytase can be defined as a phytase having a temperature-optimum of at least 60°C.

30 Preferably, the optimum temperature is determined on the

substrate phytate at pH 5.5, or on the substrate phytic acid at

10

pH 5.0. Preferred units are FYT, FTU or the units of Example 3. The phytase assay of Example 3 is most preferred.

In preferred embodiments, the optimum temperature is at least 61, 62, 63, 64, 65, 66, 67, 68, 69 or 70°C. Preferably, 5 the optimum temperature is equal to or lower than 140°C, more preferably equal to or lower than 135, 130, 125, 120, 115, 110, 105 or 100°C. Accordingly, preferred intervals of optimum temperature are: 60-140°C, 61-140°C, - (etc.) - 70-140°C; 60-135°C, 61-135°C, - (etc.) - 70-135°C; 60-130°C, - (etc.) - 70-100°C.

Preferred phytases of the present invention exhibit a degree of similarity or homology, preferably identity, to the complete amino acid sequence of either of the phytases mentioned below under (iii) - preferably to the complete amino acid sequence of Consensus-phytase-10-thermo-Q50T-K91A - of at least 48%, preferably at least 50, 52, 55, 60, 62, 65, 67, 70, 73, 75, 77, 80, 82, 85, 88, 90, 92, 95, 98 or 99%.

The degree of similarity or homology, alternatively identity, can be determined using any alignment programme known 20 in the art. A preferred alignment programme is GAP provided in the GCG version 8 program package (Program Manual for the Wisconsin Package, Version 8, August 1994, Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711) (see also Needleman, S.B. and Wunsch, C.D., (1970), Journal of Molecular Biology, 48, 443-453). Using GAP with the following settings for polypeptide sequence comparison: GAP weight of 3.000 and GAP lengthweight of 0.100.

A multiple sequence alignment can be made using the program PileUp (Program Manual for the Wisconsin Package, 30 Version 8, August 1994, Genetics Computer Group, 575 Science

Drive, Madison, Wisconsin, USA 53711), with a GapWeight of 3.000 and a GapLengthWeight of 0.100.

Using the program GAP, some selected phytases exhibit the following percentage similarity (identity in brackets) to the 5 Consensus-phytase-10-thermo(3)-Q50T-K91A amino acid sequence:

	A. fumigatus ATCC-13073 α -mutant	86.7%	(81.8%)
	Basidiomycet consensus	64.1%	(49.0%)
	Consensus-phytase-1	98.7%	(97.9%)
10	Consensus-phytase-10	96.6%	(94.4%)
	Consensus-phytase-1-thermo(8)-Q50T-K91A	97.4%	(95.5%)
	Consensus-phytase-11	96.5%	(94.2%)
	Consensus-phytase-12	92.5%	(89.9%)
	Consensus-phytase-7	95.5%	(93.4%)

A "purified" phytase is essentially free of other non-phytase polypeptides, e.g. at least about 20% pure, preferably at least about 40% pure, more preferably about 60% pure, even more preferably about 80% pure, most preferably about 90% pure, and even most preferably about 95% pure, as determined by SDS-PAGE.

Preferred thermostable phytases are the so-called consensus phytases of EP 98113176.6 (EP 0897985), viz.

- (i) any thermostable phytase which is obtainable by the processes described therein;
 - (ii) a phytase comprising the amino acid sequence shown in Fig. 2 thereof or any variant or mutein thereof, preferred muteins being those comprising the substitutions Q50L; Q50T; Q50G; Q50T-Y51N or Q50L-Y51N.

15

- (iii) a thermostable phytase which comprises at least one of the following amino acid sequence (some of which are shown in Figs. 5-12 herein), preferably the following phytases: phytase); Consensus simply Consensus-phytase-1 (or Consensus-phytase-1-thermo(3); Consensus-phytase-1-Q50T; 5 basidiomycete-consensus (or simply Basidio); Consensusphytase-10 (or Fcp 10); Consensus-phytase-11 (or Consensus Consensus-phytase-1-thermo(8)-Q50T-K91A; 11); Seq. Consensus-phytase-1-Consensus-phytase-1-thermo(8)-Q50T; Consensus-phytase-10-thermo(3)-Q50T-K91A; thermo(8); 10 Consensus-phytase-10-thermo(3)-Q50T (sometimes, "(3)" is from this expression); Aspergillus fumigatus deleted phytase lpha-mutant; Aspergillus fumigatus ATCC 13073 ATCC 13073 phytase α -mutant plus the mutations E59A, Aspergillus G404A; S364T, S126N, R329H, 15 ATCC 13073 phytase α -mutant plus the mutations E59A, K68A, Consensus-phytase-7; G404A; S364T, R329H, S126N, Consensus-phytase-12.
 - (iv) as well as thermostable variants and muteins of the phytases of (iv) and (v), in particular those comprising one or more of the following substitutions: Q50L,T,G; Q50L-Y51N; Q50T-Y51N.

The term "plant" is intended to include not only whole
25 plants as such, but also plant parts or organs, such as leaves,
seeds or grains, stem, root, tubers, flowers, callus, fruits
etc.; tissues, cells, protoplats etc.; as well as any
combinations or sub-combinations thereof. Plant tissue cultures
and plant cell lines as well as plant protoplasts are
30 specifically included herein.

The term "transgenic plant" is a plant as defined above, which has been genetically modified, as well as its progeny and propagating material thereof having retained the genetical modification. Preferably, the transgenic plant comprises at least one specific gene introduced into an ancestral plant by recombinant gene technology. The term is not confined to a single plant variety.

The invention relates to a transgenic plant which comprises a DNA-construct encoding a thermostable phytase.

In a preferred embodiment the transgenic plant is a plant grouping which is characterized in that it comprises a DNA-construct encoding a thermostable phytase. The members of this plant grouping may very well possess individuality, but are clearly distinguishable from other varieties by their common characteristic feature of the the thermostable phytase DNA-construct.

Accordingly, the present teaching is applicable to more than one plant variety. No naturally occurring plant varieties are included amongst the plants of the invention.

In another preferred embodiment the invention relates to a transgenic plant variety or a variant thereof; a transgenic plant species, a transgenic plant genus, a transgenic plant family, and/or a transgenic plant order. More preferably, plant varieties as such are disclaimed.

Any thermostable phytase may be used in the present invention, e.g. any wild-type phytases, genetically engineered phytases, consensus phytases, phytase muteins, and/or phytase variants. Genetically engineered phytases include, but are not limited to, phytases prepared by site-directed mutagenesis, gene shuffling, random mutagenesis, etc.

14

The nucleotide sequence encoding a wild-type thermostable phytase may be of any origin, including mammalian, plant and microbial origin and may be isolated from these sources by conventional methods. Preferably, the nucleotide sequence is derived from a microorganism, such as a fungus, e.g. a yeast or a filamentous fungus, or a bacterium. The DNA sequence encoding a thermostable phytase may be isolated from the cell producing it, using various methods well known in the art (see e.g. WO 98/28409 and EP 0897985).

The nucleotide sequence encoding a thermostable genetically engineered or consensus phytase, including muteins and variants thereof, may be prepared in any way, e.g. as described in Example 3 hereof and in EP 0897985.

In order to accomplish expression of the thermostable phytase in a plant of the invention the nucleotide sequence encoding the phytase is inserted into an expression construct containing regulatory elements or sequences capable of directing the expression of the nucleotide sequence and, if necessary or desired, to direct secretion of the gene product or targetting of the gene product to the seeds of the plant.

In order for transcription to occur the nucleotide sequence encoding the thermostable phytase is operably linked to a suitable promoter capable of mediating transcription in the plant in question. The promoter may be an inducible promoter or a constitutive promoter. Typically, an inducible promoter mediates transcription in a tissue-specific or growth-stage specific manner, whereas a constitutive promoter provides for sustained transcription in all cell tissues. An example of a suitable constitutive promoter useful for the present invention is the cauliflower mosaic virus 35 S promoter. Transcription initiation sequences from the tumor-inducing plasmid (Ti) of

Agrobacterium such as the octopine synthase, nopaline synthase, or mannopine synthase initiator, are further examples of preferred constitutive promoters.

Examples of suitable inducible promoters include a seed5 specific promoter such as the promoter expressing alpha-amylase in wheat seeds (see Stefanov et al, Acta Biologica Hungarica. Vol. 42, No. 4 pp. 323-330 (1991), a promoter of the gene encoding a rice seed storage protein such as glutelin, prolamin, globulin or albumin (Wu et al., Plant and Cell Physiology Vol. 39, No. 8 pp. 885-889 (1998)), a Vicia faba promoter from the legumin B4 and the unknown seed protein gene from Vicia faba described by Conrad U. et al, Journal of Plant Physiology Vol. 152, No. 6 pp. 708-711 (1998), the storage protein napA promoter from Brassica napus, or any other seed specific promoter known in the art, eg as described in WO 91/14772.

In order to increase the expression of the thermostable phytase it is desirable that a promoter enhancer element is used. For instance, the promoter enhancer may be an intron which is placed between the promoter and the amylase gene. The intron 20 may be one derived from a monocot or a dicot. For instance, the intron may be the first intron from the rice Waxy (Wx) gene (Li et al., Plant Science Vol. 108, No. 2, pp. 181-190 (1995)), the first intron from the maize Ubil (Ubiquitin) gene (Vain et al., Plant Cell Reports Vol. 15, No. 7 pp. 489-494 (1996)) or the first intron from the Actl (actin) gene. As an example of a dicot intron the chsA intron (Vain et al. op cit.) is mentioned. Also, a seed specific enhancer may be used for increasing the expression of the thermostable phytase in seeds. An example of a seed specific enhancer is the one derived from the beta-

(Phaseolus vulgaris) disclosed by Vandergeest and Hall, Plant Molecular Biology Vol. 32, No. 4, pp. 579-588 (1996).

Also, the expression construct preferably contains a terminator sequence to signal transcription termination of the the thermostable phytase gene such as the rbcS2' and the nos3' terminators.

successfully transformed facilitate selection of plants, the expression construct should also preferably include one or more selectable markers, e.g. an antibiotic resistance 10 selection marker or a selection marker providing resistance to a herbicide. One widely used selection marker is the neomycin which provides kanamycin gene (NPTII) phosphotransferase resistance. Examples of other suitable markers include a marker providing a measurable enzyme activity, e.g. dihydrofolate b-glucoronidase luciferase, and 15 reductase, Phosphinothricin acetyl transferase may be used as a selection marker in combination with the herbicide basta or bialaphos.

The transgenic plant of the invention may be prepared by methods known in the art. The transformation method used will 20 depend on the plant species to be transformed and can be selected from any of the transformation methods known in the art such as Agrobacterium mediated transformation (Zambryski et al., EMBO Journal 2, pp 2143-2150, 1993), particle bombardment, electroporation (Fromm et al. 1986, Nature 319, pp 791-793), and 25 virus mediated transformation. For transformation of monocots bombardment (ie biolistic transformation) of particle embryogenic cell lines or cultured embryos are preferred. Below, references are listed, which disclose various methods 1991, al. transforming various plants: Rice (Cristou et 30 Bio/Technology 9, pp. 957-962), Maize (Gordon-Kamm et al. 1990, 1992, pp. 603-618), Oat (Somers al. et Plant Cell 2,

Bio/Technology 10, pp 1589-1594), Wheat (Vasil et al. 1991,
Bio/Technology 10, pp. 667-674, Weeks et al. 1993, Plant
Physiology 102, pp. 1077-1084) and Barley (Wan and Lemaux 1994,
Plant Physiology 102, pp. 37-48, review Vasil 1994, Plant Mol.
5 Biol. 25, pp 925-937).

More specifically, Agrobacterium mediated transformation is conveniently achieved as follows:

A vector system carrying the thermostable phytase is constructed. The vector system may comprise of one vector, but it can comprise of two vectors. In the case of two vectors the vector system is referred to as a binary vector system (Gynheung An et al.(1980), Binary Vectors, Plant Molecular Biology Manual A3, 1-19).

plant transformation vector Agrobacterium based An both E.coli replication origin(s) for and of 15 consists Agrobacterium and a bacterial selection marker. A right and preferably also a left border from the Ti plasmid plasmid Ri from the Agrobacterium tumefaciens or Agrobacterium rhizogenes is nessesary for the transformation of 20 the plant. Between the borders the expression construct is placed which contains the thermostable phytase gene appropriate regulatory sequences such as promotor and terminator sequences. Additionally, a selection gene e.g. the neomycin phosphotransferase type II (NPTII) gene from transposon Tn5 and 25 a reporter gene such as the GUS (betha-glucuronidase) gene is cloned between the borders. A disarmed Agrobacterium strain harboring a helper plasmid containing the virulens genes is transformed with the above vector. The transformed Agrobacterium strain is then used for plant transformation.

The invention also relates to a method of preparing a transgenic plant capable of expressing a thermostable phytase,

said method comprising the steps of (i) isolating a nucleotide sequence encoding a thermostable phytase; (ii) inserting the nucleotide sequence of (i) in an expression construct capable of mediating the expression of the nucleotide sequence in a selected host plant; and (iii) transforming the selected host plant with the expression construct.

The above method in which "at least one" replaces "a," when used in relation to the thermostable phytase, is also within this invention.

This method is an essentially non-biological method.

Any plant may be a selected host plant. More specifically, the plant can be dicotyledonous or monocotyledonous, for short a dicot or a monocot. Of primary interest are such plants which are potential food or feed components. These plants may comprise phytic acid. Examples of monocot plants are grasses, such as meadow grass (blue grass, Poa), forage grass such as festuca, lolium, temperate grass, such as Agrostis, and cereals, e.g. wheat, oats, rye, barley, rice, sorghum and maize (corn).

Examples of dicot plants are legumes, such as lupins, pea, 20 bean and soybean, and cruciferous (family Brassicaceae), such as cauliflower, oil seed rape and the closely related model organism Arabidopsis thaliana.

Of particular interest are monocotyledonous plants, in particular crops or cereal plants such as wheat (Triticum, e.g. 25 aestivum), barley (Hardeum, e.g. vulgare), oats, rye, rice, sorghum and corn (Zea, e.g. mays).

Of further particular interest are dicotyledonous plants, such as those mentioned above.

In a preferred embodiment, the ancestral plant or host 30 plant is per se a desired feed ingredient.

Examples

Example 1

FYT-assay - for analyzing phytase enzyme preparations

The phytase activity can be measured using the following assay:

- 5 10 μl diluted enzyme samples (diluted in 0.1 M sodium acetate, 0.01 % Tween20, pH 5.5) are added into 250 μl 5 mM sodium phytate (Sigma) in 0.1 M sodium acetate, 0.01 % Tween20, pH 5.5 (pH adjusted after dissolving the sodium phytate; the substrate is preheated) and incubated for 30 minutes at 37°C. The reaction
- 10 is stopped by adding 250 μl 10 % TCA and free phosphate is measured by adding 500 μl 7.3 g FeSO4 in 100 ml molybdate reagent (2.5 g (NH₄)₆Mo₇O₂₄.4H₂O in 8 ml H₂SO₄ diluted to 250 ml). The absorbance at 750 nm is measured on 200 μl samples in 96 well microtiter plates. Substrate and enzyme blanks are included. A phosphate standard curve is also included (0-2 mM phosphate). 1 FYT equals the amount of enzyme that releases 1
 - phosphate). 1 FYT equals the amount of enzyme that releases 1 µmol phosphate/min at the given conditions. This assay is preferred for phytase enzyme preparations (when not in admixture with other feed ingredients).

20

Example 2

FTU assay - for analyzing phytase in admixture with feed ingredients

One FTU is defined as the amount of enzym, which at stan25 dard conditions (37°C, pH 5,5; reaction time 60 minutes and
start concentration of phytic acid 5 mM) releases phosphate
equivalent to 1 µmol phosphate per minute.

1 FTU = 1 FYT

The FTU assay is preferred for phytase activity measure-30 ments on animal feed premixes and the like complex compositions.

Reagents /substrates

Extraction buffer for feed etc.

This buffer is also used for preparation of PO_4 -standards and further dilution of premix samples.

5 0.22 M acetate buffer with Tween 20 pH 5.5

30 g sodium acetate trihydrate (MW = 136,08 g/mol) e.g. Merck Art 46267 per liter and 0,1 g Tween 20 e.g. Merck Art 22184 pr. liter are weighed out.

The sodium acetate is dissolved in demineralised water.

Tween 20 is added, and pH adjusted to $5,50 \pm 0,05$ with acetic acid.

Add demineralised water to total volume.

Extraction buffer for premix

- 0,22 M acetate buffer with Tween 20, EDTA, PO_4^{3-} og BSA.
- 15 30 g sodium acetate trihydrate e.g. Merck Art 6267 per liter.
 - 0,1 g Tween 20 e.g. Merck Art 22184 per liter.
 - 30 g EDTA f.eks. Merck Art 8418 pr. liter.
 - 20 g Na₂HPO₄, 2H₂O e.g. Merck Art 6580 per liter.
- 20 0,5 g BSA (Bovine Serum Albumine, e.g. Sigma Art A-9647 per liter.

The ingredients are dissolved in demineralised water, and pH is adjusted to 5.50 \pm 0.05 with acetic acid.

Add demineralised water to total volume.

25 BSA is not stable, and must therefore be added the same day the buffer is used.

50 mM PO43-stock solution

0,681 g KH2PO4 (MW = 136,09 g/mol) e.g. Merck Art 4873 is weighed out and dissolved in 100 ml 0,22 M sodium acetat with Tween, pH 5,5.

5 Storage stability: 1 week in refrigerator.

0.22 M acetate buffer pH 5.5 without Tween

This buffer is used for production of phytic acid substrate).

150 g sodium acetate trihydrate (MW = 136,08) e.g. Merck 10 Art 6267 is weighed out and dissolved in demineralised water, and pH is adjusted with acetic acid to $5,50 \pm 0,05$.

Add demineralised water to 5000 ml.

Storage stability: 1 week at room temperature.

Phytic acid substrate: 5 mM phytic acid

The volume of phytic acid is calculated with allowance for the water content of the used batch.

If the water content is e.g. 8,4 % the following is obtained:

20
$$\frac{0,005 \, mol \, / \, l \times 923,8 \, g \, / \, mol}{\left(1 \div 0,084\right)} = 5,04 \, g \, / \, l$$

Phytic acid (Na-salt) (MW = 923,8 g/mol) e.g. Sigma P-8810 is weighed out and dissolved in 0,22 M acetate buffer (without tween). Addition of (diluted) acetic acid increases the dissolution speed.

pH is adjusted to 5,50 \pm 0,05 with acetic acid. Add 0,22 M acetate buffer to total volume.

21.7 % nitric acid solution

For stop solution.

1 part concentrated (65%) nitric acid is mixed into 2 parts demineralised water.

Molybdate reagent

5 For stop solution.

100 g ammonium heptamolybdate tetrahydrate (NH $_4$) 6Mo $_7$ O $_{24}$, 4H $_2$ O e.g. Merck Art 1182 is dissolved in demineralised water. 10 ml 25 % NH $_3$ is added.

Add demineralised water to 1 liter.

10 0.24 % Ammonium vanadate

Bought from fra Bie & Berntsen.

Molybdat/vanadat stop solution

1 part vanadate solution (0,24 % ammonium vanadate) + 1 part molybdate solution are mixed. 2 parts 21,7 % nitric acid solution are added.

The solution is prepared not more than 2 hours before use, and the bottle is wrapped in tinfoil.

Samples

Frozen samples are defrosted in a refrigerator overnight.

20 Sample size for feed samples: At least 70 g, preferably 100 g.

Feed samples

Choose a solution volume which allows addition of buffer corresponding to 10 times the sample weight, e.g. 100 g is dissolved in 1000 ml 0,22 M acetate buffer with Tween, see enclosure 1. Round up to nearest solution volume.

If the sample size is approx. 100 g all the sample is ground in a coffee grinder and subsequently placed in tared

beakers. The sample weight is noted. It is not necessary to grind not-pelleted samples. If a sample is too big to handle, it is sample split into parts of approx. 100 g.

Magnets are placed in the beakers and 0,22 M acetate 5 buffer with Tween is added.

The samples are extracted for 90 minutes.

After extraction the samples rest for 30 minuts to allow for the feed to sediment. A 5 ml sample is withdrawn with a pipette. The sample is taken 2 - 5 cm under the surface of the solution and placed in a centrifuge glass, which is covered by a lid.

The samples are centrifuged for 10 minutes at 4000 rpm.

Premix samples

Choose a solution volume which allows addition of buffer 15 corresponding to 10 times the sample weight. Round up to nearest solution volume.

If the samples have been weighed (50 - 100 g) all of the sample is placed in tared beakers. The sample weight is noted. If a sample is too big to handle, it is split into parts of ap-20 prox. 100 g.

Magnets are placed in the beakers and 0,22 M acetate buffer with Tween, EDTA og PO_4^{3-} is added.

The samples are extracted for 60 minutes.

After extraction the samples rest for 30 minutes to allow for the premix to sediment. A 5 ml sample is withdrawn with a pipette. The sample is taken 2 - 5 cm under the surface of the solution and placed in a centrifuge glass, which is covered by a lid.

The samples are centrifuged for 10 minutes at 4000 rpm.

Analysis

Extracts of feed samples are analysed directly.

Extracts of premix are diluted to approx. 1,5 FTU/g (A_{415} 5 (main sample) < 1,0).

0,22 M acetate buffer with Tween 20 is used for the dilution.

Main Samples

 2×100 ml of the supernatant from the extracted and centrifuged samples are placed in marked glass test tubes and a magnet is placed in each tube.

When all samples are ready they are placed on a water bath with stirring. Temperature: 37 °C.

3,0 ml substrate is added.

Incubation for exactly 60 minutes after addition of substrate.

The samples are taken off the water bath and 2,0 ml stop solution is added (exactly 60 minutes after addition of substrate).

The samples are stirred for 1 minute or longer.

Feed samples are centrifuged for 10 minutes at 4000 rpm (It is not necessary to centrifuge premix samples).

Blind samples

100 ml of the supernatant from the extracted and centri-25 fuged samples are placed in marked glass test tubes, and a magnet is placed in each tube.

2,0 ml stop solution is added to the samples.

3,0 ml substrate is added to the samples.

The samples are incubated for 60 minutes at room temperature.

The feed samples are centrifuged for 10 minutes at 4000 5 rpm (it is not necessary to centrifuge premix samples).

Standards

 2×100 ml are taken from each of the 8 standards and also 4×100 ml 0,22 M acetate buffer (reagent blind).

 A_{415} is measured on all samples.

10 CALCULATION

 $FTU/g = \mu mol PO_4^{3-} / (min * g (sample))$

C g sample is weighed out (after grinding).

15 100 μ l is taken from the extracted and centrifuged sample.

PO,3- standard curve is linear.

From the regression curve for the PO_4^{3-} standard the actual con-20 centration of the sample is found (concentration in mM):

 $[PO_4^{3-}] = (x - b) / a$ $x = A_{415}$ a = slope b = intercept with y-axis

25 μ mol PO₄³⁻/min = { [PO₄³⁻] (mM) × Vol (liter) × 1000 μ mol/mmol } /t

t = incubation time in minutes.

Vol = sample volume in liter = 0,0001 liter

1000 = conversion factor from mmol to μ mol

FTU
$$/g_{pxeve} = \{ (x - b) \times Vol \times 1000 \times F_p \} / \{ a \times t \times C \}$$

C = gram sample weighed out

5 F_P = Relation between the sample taken out and the total sample (after extraction). Example: 0,100 ml taken from 1000 ml \rightarrow F_P = 1000/0,100 = 10000.

Reduced expression with insertion of the following values:

10 t = 60

Vol = 0,0001 l

 $F_{\rm P} = 10000$

FTU $/g_{\text{sample}} = \{ (x - b) \times 0,0001 \times 1000 \times 10000 \} / \{ a \times 60 \times C \}$

15 Example 3

Determination of optimum temperature and melting point Tm of various phytases

The thermostability of various phytases has been determined, viz. the melting temperature, Tm, and/or the optimum temperature.

The phytase of Aspergillus niger NRRL 3135 was prepared as described in EP 0420358 and van Hartingsveldt et al (Gene, 127, 87-94, 1993).

The phytases of Aspergillus fumigatus ATCC 13073, 25 Aspergillus terreus 9A-1, Aspergillus terreus CBS 116.46, Aspergillus nidulans, Myceliophthora thermophila, and Talaromyces thermophilus were prepared as described in EP-0897985 and the references therein.

Consensus-phytase-1 (Fig. 5) and Consensus-phytase-1-Q50T are shown in and were prepared as described in EP 0897985.

Consensus-phytase-10 was derived and prepared according to the teachings of EP-0897985 (Examples 1-2 and 3-7.respectively), however adding to the alignment at Fig. 1 thereof the phytase sequence of Thermomyces lanuginosa (Berka et al, 5 Appl. Environ. Microbiol. 64, 4423-4427, 1998) and basidiomycete consensus sequence (derivation described below), omitting the sequence of A.niger T213, and assigning a vote weight of 0.5 for the remaining A.niger phytase sequences. The derivation of the sequence of Consensus-phytase-10 is shown in 10 Fig. 7.

The basidiomycete consensus sequence was also derived according to the principles of EP-0897985, viz. from the five basidiomycete phytases of WO 98/28409, starting with the first amino acid residue of the mature phytases (excluding signal peptide). A vote weight of 0.5 was assigned to the two Paxillus phytases, all other genes were used with a vote weight of 1.0 - see Fig. 6.

The muteins Consensus-phytase-10-thermo, Consensus-phytase-10-thermo-Q50T-K91A (Fig. 10) and Consensus-phytase-10-20 thermo-Q50T were prepared from consensus-phytase-10, in analogy to Examples 5-8 of EP-0897985, by introducing the three back-mutations K94A, V158I and A396S ("thermo(3)" or "thermo") and, where applicable, also the mutations Q50T or Q50T-K91A.

The muteins Consensus-phytase-1-thermo(8), Consensus25 phytase-1-thermo(8)-Q50T-K91A (Fig. 9) and Consensus-phytase-1thermo(8)-Q50T, were prepared from consensus-phytase-1, in
analogy to Example 8 of EP-0897985, by introducing the eight
mutations E58A, D197N, E267D, R291I, R329H, S364T, A379K and
G404A ("thermo(8)") and, where applicable, also the mutations
30 Q50T or Q50T-K91A.

Consensus-phytase-1-thermo(3) was prepared from consensus-phytase-1 by introduction of the three mutations K94A, V158I and A396S.

An Aspergillus fumigatus so-called α -mutant (with the mutations Q51(27)T, F55Y, V100I, F114Y, A243L, S265P, N294D) and the further muteins thereof shown in Table 1 were prepared asgenerally described above. The position numbering refers to Fig. 11 hereof, except for the number in parentheses which refers to the numbering used in EP 0897010.

DNA constructs encoding the above thermostable phytases can be prepared e.g. according to the teachings of EP 0897985. For expression thereof in plants, reference is made to the present description.

In order to determine the unfolding temperature or melting temperature, Tm, of a phytase, differential scanning calorimetry was applied as previously published by Brugger et al (1997): "Thermal denaturation of phytases and pH 2.5 acid phosphatase studied by differential scanning calorimetry," in The Biochemistry of phytate and phytase (eds. Rasmussen, S.K; Raboy, V.; Dalbøge, H. and Loewus, F.; Kluwer Academic Publishers).

Homogenous or purified phytase solutions of 50-60 mg/ml of protein are prepared, and extensively dialyzed against 10 mM sodium acetate, pH 5.0. A constant heating rate of 10°C/min is applied up to 90-95°C.

The results of Tm determinations on the above phytases are shown in Table 1 below; for selected phytases also in Figs. 1-4.

In Table 1 below, the optimum temperature of various phytases is also indicated. For this determination, phytase activity was determined basically as described by Mitchell et al (Microbiology 143, 245-252, 1997): The activity was measured in an assay mixture containing 0.5% phytic acid (~ 5 mM) in 200 mM

25

sodium acetate, pH 5.0. After 15 min of incubation at 37°C, the reaction was stopped by addition of an equal volume of 15% trichloroacetic acid. The liberated phosphate was quantified by mixing 100 µl of the assay mixture with 900µl H₂O and 1 ml of 5 0.6 M H₂SO₄, 2% ascorbic acid and 0.5% ammonium molybdate. Standard solutions of potassium phosphate were used asreference. One unit of enzyme activity was defined as the amount of enzyme that releases 1 µmol phosphate per minute at 37°C. The protein concentration was determined using the enzyme extinction coefficient at 280 nm calculated according to Pace et al (Prot.Sci. 4, 2411-2423, 1995): Consensus phytase, 1.101; consensus phytase 7, 1.068; consensus phytase 10, 1.039.

For determination of the temperature optimum, enzyme (100µl) and substrate solution (100µl) were pre-incubated for 5 min at the given temperature. The reaction was started by addition of the substrate solution to the enzyme. After 15 min incubation, the reaction was stopped with trichloroacetic acid and the amount of phosphate released was determined. Phytase-activity-versus-temperature is plotted, and the temperature optimum is determined as that temperature at which the acitivity reaches its maximum value.

Table 1
Temperature optimum and Tm for various phytases

Phytase Optimum temperature (°C)

Aspergillus niger 55 63.3

NRRL 3135

Aspergillus 55 62.5

fumigatus ATCC 13073

31

S126N, R329H, S364T, G404A		
Aspergillus	63	-
fumigatus — as		
above, plus mutation		
K68A		-
Aspergillus	60	67.0
fumigatus α-mutant		
(Q51(27)T, F55Y,		
V100I, F114Y, A243L,		
S265P, N294D)		

CLAIMS

25

- A process of preparing an animal feed, which process comprises an agglomeration of feed ingredients, wherein a thermostable phytase is added before or during the sagglomeration.
 - 2. The process of claim 1, wherein the feed ingredients are heated to a temperature of at least 65°C.
- 10 3. The process of any of claims 1-2, wherein the thermostable phytase is a phytase with a Tm as measured by DSC of at least 65°C, using for the DSC a constant heating rate of 10°C/min.
- 4. The process of any of claims 1-3, when performed in a feed 15 expander.
 - 5. The process of any of claims 1-3, when performed in an extruder.
- 20 6. The process of any of claims 1-3, when performed in a pellet press.
 - 7. The process of any of claims 1-6, wherein the thermostable phytase is present in a transgenic plant.
 - 8. The process of any of claims 1-7, wherein the agglomeration includes the following steps:
 - (a) pre-heating the feed ingredients to a temperature of at least 45°C; and
- 30 (b) heating the product of step (a) to a temperature of at least 65°C;


33


wherein the thermostable phytase is added prior to or during step (a) and/or (b).


- 9. A transgenic plant which comprises a DNA-construct 5 encoding a thermostable phytase.
- 10. The transgenic plant of claim 9, wherein the DNA-construct encoding the thermostable phytase is operably linked to regulatory sequences capable of mediating expression of said phytase encoding sequence in at least one part of the plant.
- 11. An expression construct which comprises a DNA construct encoding a thermostable phytase, operably linked to regulatory sequences capable of mediating expression of said phytase encoding sequence in at least one part of a plant.
 - 12. A vector which comprises the expression construct of claim 11.
- 20 13. A method of preparing a transgenic plant capable of expressing a thermostable phytase, said method comprising the steps of
 - (i) isolating a nucleotide sequence encoding a thermostable phytase;
- 25 (ii) inserting the nucleotide sequence of (i) in an expression construct capable of mediating the expression of the nucleotide sequence in a selected host plant; and
 - (iii) transforming the selected host plant with the expression construct.


34


14. The method of claim 13, which comprises the further step of extracting the phytase from the plant.














KhsDCNSVDh GYQCFPELSH kWGlYAPYFS LQDESPFPlD VPEDChITFV A. terreus 9A-1 MhsDCTSVDr GYQCFPELSH kWGlYAPYFS LQDESPFPlD VPDDChITFV A. terreus cbs A. niger var. awamori NqsTCDTVDQ GYQCFSETSH LWGQYAPFFS LANESAISPD VPAGCrVTFA NGSSCDTVDQ GYQCFSETSH LWGQYAPFFS LANESVISPD VPAGCTVTFA A. niger T213 NGSSCDTVDQ GYQCFSETSH LWGQYAPFFS LANESVISPE VPAGCIVTFA A. niger NRRL3135 GSkSCDTVD1 GYQCsPATSH LWGQYSPFFS LEDE1SVSSK LPKDCrITLV A. fumigatus 13073 A. fumigatus 32722 GSKSCDTVDl GYQCsPATSH LWGQYSPFFS LEDELSVSSK LPKDCrITLV GSKSCDTVD1 GYQCsPATSH LWGQYSPFFS LEDE1SVSSK LPKDCrITLV A. fumigatus 58128 GSKSCDTVD1 GYQCsPATSH LWGQYSPFFS LEDELSVSSK LPKDCrITLV A. fumigatus.26906 GS%ACDTVE1 GYQCsPGTSH LWGQYSPFFS LEDE1SVSSD LPKDCTVTFV A. fumigatus 32239 ONHSCNTADG GYQCFPNVSH VWGQYSPYFS IEQESAISED VPHGCeVTFV E. nidulans DSHSCNTVEG GYQCTPEISH SWGQYSPFFS LADQSEISPD VPQNCKITFV T. thermophilus ESRPCDTpDl GFQCgTAISH FWGQYSPYFS VpSElDaS.. IPDDCeVTFA M. thermophila NSHSCDTVDG GYQCFPEISH LWGQYSPYFS LEDESAISPD VPDDC-VTFV Consensus NSHSCDTVDG GYQCFPEISH LWGQYSPYFS LEDESAISPD VPDDCRVTFV Consensus phytase 100 QVLARHGARS PThSKtKAYA AtlAAlQKSA TaFpGKYAFL QSYNYSLDSE A. terreus 9A-1 QVLARHGARS PTDSKtKAYA AtlaalQKNA TalpGKYAFL KSYNYSMGSE A. terreus cbs A. niger var. awamori QVLSRHGARY PTESKGKKYS ALIEEIQQNV TtFDGKYAFL KTYNYSLGAD QVLSRHGARY PTESKGKKYS ALIEEIQQNV TtFDGKYAFL KTYNYSLGAD A. niger T213 QVLSRHGARY PTDSKgKkYS ALIEEIQQNA TtFDGKYAFL KTYNYSLGAD A. niger NRRL3135 QVLSRHGARY PTSSKskkyk klvtaiqana tdfkgkfafl ktynytlgad A. fumigatus 13073 QVLSRHGARY PTSSKsKkYK kLVTAIQaNA TdFKGKFAFL KTYNYTLGAD A. fumigatus 32722 QVLSRHGARY PTSSKsKkYK kLVTAIQANA TdFKGKFAFL KTYNYTLGAD A. fumigatus 58128 QVLSRHGARY PTSSKsKkYK kLVTAIQaNA TdFKGKFAFL KTYNYTLGAD A. fumigatus 26906 OVLSRHGARY PTASKSKKYK KLVTAIQKNA TEFKGKFAFL ETYNYTLGAD A. fumigatus 32239 QVLSRHGARY PTESKSKAYS GLIEAIQKNA TSFWGQYAFL ESYNYTLGAD E. nidulans QLLSRHGARY PTSSKtElys QLISTIQKTA TayKGyYAFL KDYTYqLGAN T. thermophilus QVLSRHGARa PT1KRaaSYv DLIDrIHhGA ISYGPGYEFL RTYDYTLGAD M. thermophila QVLSRHGARY PTSSK-KAYS ALIEAIQKNA T-FKGKYAFL KTYNYTLGAD Consensus QVLSRHGARY PTSSKSKAYS ALIEAIQKNA TAFKGKYAFL KTYNYTLGAD Consensus phytase 101 150 ELTPFGrNQL rDlGaQFYeR YNALTRhinP FVRATDASRV hESAEKFVEG A. terreus 9A-1 NLTPFGrNQL qDlGaQFYRR YDTLTRhinP FVRAADSSRV hESAEKFVEG A. terreus cbs A. niger var. awamori DLTPFGEQEL VNSGIKFYQR YESLTRNIIP FIRSSGSSRV IASGEKFIEG DLTPFGEQEL VNSGIKFYQR YESLTRNIIP FIRSSGSSRV IASGEKFIEG A. niger T213 DLTPFGEQEL VNSGIKFYQR YESLTRNIVP FIRSSGSSRV IASGKKFIEG A. niger NRRL3135 DLTPFGEQQL VNSGIKFYQR YKALARSVVP FIRASGSDRV IASGEKFIEG A. fumigatus 13073 DLTPFGEQQL VNSGIKFYQR YKALARSVVP FIRASGSDRV IASGEKFIEG A. fumigatus 32722 DLTPFGEQQL VNSGIKFYQR YKALARSVVP FIRASGSDRV IASGEKFIEG A. fumigatus 58128 DLTAFGEQQL VNSGIKFYQR YKALARSVVP FIRASGSDRV IASGEKFIEG A. fumigatus 26906 DLTPFGEQQM VNSGIKFYQK YKALAGSVVP FIRSSGSDRV IASGEKFIEG A. fumigatus 32239 DLTIFGENOM VDSGaKFYRR YKNLARKnTP FIRASGSDRV VASAEKFING E. nidulans DLTPFGENQM IQ1GIKFYnH YKSLARNAVP FVRCSGSDRV IASGrlFIEG T. thermophilus ELTREGQQQM VNSGIKFYRR YRALARKSIP FVRTAGQDRV VhSAENFTQG M. thermophila DLTPFGENQM VNSGIKFYRR YKALARK-VP FVRASGSDRV IASAEKFIEG Consensus DLTPFGENOM VNSGIKFYRR YKALARKIVP FIRASGSDRV IASAEKFIEG Consensus phytase

PCT/DK99/00154 WO 99/48380

151

Consensus phytase

6/32

FQTARqDDHh ANPHQPSPrV DVaipegsay NNTLEHSlCT AFES...STV 200 FQNARQGDPh ANPHQPSPIV DVVIPEGTAY NNTLEHSICT AFEA...STV A. terreus 9A-1 A. niger var. awamori FQSTKLkDPr AqpgQSSPkI DVVISEASSS NNTLDPGTCT VFED...SEL FQSTKLkDPr AqpgQSSPkI DVVISEASSS NNTLDPGTCT VFED...SEL FQSTKLkDPr AqpgQSSPkI DVVISEASSB NNTLDPGTCT VFED...SEL A. niger T213 A. niger NRRL3135 FQQAKLADPG A.TNRAAPAI SVIIPESETF NNTLDHGVCT kFEA...SQL A. fumigatus 13073 FQQAKLADPG A.TNRAAPAI SVIIPESETF NNTLDHGVCT kFEA...SQL A. fumigatus 32722 FQQAKLADPG A.TNRAAPAI SVIIPESETF NNTLDHGVCT kFEA...SQL A. fumigatus 58128 FQQAKLADPG A.TNRAAPAI SVIIPESETF NNTLDHGVCT kFEA...SQL A. fumigatus 26906 FQQANVADPG A.TNRAAPVI SVIIPESETY NNTLDHSVCT NFEA...SEL FRKAQLADHG S... GQATPVV NVIIPEIDGF NNTLDHSTCV SFEN...DEr A. fumigatus 32239 FQSAKVlDPh SDkHDAPPTI NVIIEEGPSY NNTLDtGSCP VFED...SSg E. nidulans T. thermophilus FHSALLADRG STVRPTIPYS MVVIPETAGA NNTLHNDLCT AFEEgpySTI M. thermophila FQSAKLADPG S-PHQASPVI NVIIPEGSGY NNTLDHGTCT AFED---SEL FQSAKLADPG SQPHQASPVI DVIIPEGSGY NNTLDHGTCT AFED...SEL Consensus Consensus phytase 201 GDDAVANFTA VFAPAIaQRL EADLPGVqLS TDDVVnLMAM CPFETVSlTD 250 GDAAADNFTA VFAPAIAKRL EADLPGVQLS ADDVVnLMAM CPFETVSlTD A. terreus 9A-1 A. niger var. awamori ADTVEANFTA TFAPSIRQRL ENDLSGVTLT DTEVTYLMDM CSFDTIStST A. niger T213 ADTVEANFTA TFAPSIRORL ENDLSGVTLT DTEVTYLMDM CSFDTIStST ADTVEANFTA TFVPSIRQRL ENDLSGVTLT DTEVTYLMDM CSFDTISLST GDEVAANFTA 1FAPDIRARA EKHLPGVTLT DEDVV8LMDM CSFDTVARTS A. niger NRRL3135 A. fumigatus 13073 GDEVAANFTA 1FAPDIRARA EKHLPGVTLT DEDVVSLMDM CSFDTVARTS A. fumigatus 32722 A. fumigatus 58128 GDEVAANFTA 1FAPDIRARA EKHLPGVTLT DEDVV8LMDM CSFDTVARTS GDEVAANFTA 1FAPDIRARA KKHLPGVTLT DEDVVSLMDM CSFDTVARTS A. fumigatus 26906 GDEVEANFTA 1FAPAIRARI EKHLPGVQLT DDDVVSLMDM CSFDTVARTA A. fumigatus 32239 ADEIEANFTA IMGPPIRKRL ENDLPGIKLT NENVIYLMDM CSFDTMARTA GHDAQEKFAK QFAPAILEKI KDHLPGVDLA VSDVPYLMDL CPFETLARNH E. nidulans T. thermophilus GDDAQDTYLS TFAGPITARV NANLPGANLT DADTVALMOL CPFETVASSS M. thermophila GDDAEANFTA TFAPAIRARL EADLPGVTLT DEDVV-LMDM CPFETVARTS GDDVEANFTA LFAPAIRARL EADLPGVTLT DEDVVYLMDM CPFETVARTS Consensus Consensus phytase 251DAhTLSPFC DLFTAtEWtq YNYL1SLDKY YGYGGGNPLG 300 DAhTLSPFC DLFTAREWLQ YNYLlSLDKY YGYGGGNPLG A. terreus 9A-1 A. niger var. awamorivDTKLSPFC DLFTHdEWih YDYLQSLkKY YGHGAGNPLG A. niger T213 A. niger NRRL3135DASQLSPFC QLFTHnEWkk YNYLQSLGKY YGYGAGNPLG A. fumigatus 13073 A. fumigatus 32722DASQLSPFC QLFTHnEWkk YNYLQSLGKY YGYGAGNPLG A. fumigatus 58128 A. fumigatus 26906DASELSPFC AIFTHDEWAK YDYLQSLGKY YGYGAGNPLG A. fumigatus 32239 E. nidulans sdpatadagg gNGrpLSPFC rLFSEsEWra YDYLQSVGKW YGYGPGNPLG T. thermophilus M. thermophila Consensus

A. fumigatus 32239

E. nidulans

7/32

301 350 PVQCVGWaNE LMARLTRAPV HDHTCVNNTL DASPATFPLN ATLYADFSHD A. terreus 9A-1 PVQGVGWANE LIARLTRSPV HDHTCVNNTL DANPATFPLN ATLYADFSHD A. terreus cbs A. niger var. awamori PTQGVGYANE LIARLTHSPV HDDTSSNHTL DSNPATFPLN STLYADFSHD PTQCVGYANE LIARLTHSPV HDDTSSNHTL DSNPATFPLN STLYADFSHD A. niger T213 PTQGVGYANE LIARLTHSPV HDDTSSNHTL DSSPATFPLN STLYADFSHD A. niger NRRL3135 A. fumigatus 13073 PAQGIGFTNE LIARLTRSPV QDHTSTNSTL VSNPATFPLN ATMYVDFSHD PAQGIGFUNE LIARLTRSPV QDHTSTNSTL VSNPATFPLN ATMYVDFSHD A. fumigatus 32722 A. fumigatus 58128 A. fumigatus 26906 PAQGIGFTNE LIARLTRSPV QDHTSTNsTL VSNPATFPLN ATMYVDFSHD PAQGIGFTNE LIARLTRSPV QDHTSTNsTL VSNPATFPLN ATMYVDFSHD PAQGIGFTNE LIARLINSPV QDHTSTNSTL DSDPATFPLN ATIYVDFSHD A. fumigatus 32239 PAQGIGFTNE LIARLTQSPV QDNTSTNHTL DSNPATFPLD rKLYADFSHD E. nidulans PAQGVGFVNE LIARMTHSPV QDYTTVNHTL DSNPATFPLN ATLYADFSHD T. thermophilus PTQGVGFvNE LLARLAGVPV RDgTSTNRTL DGDPrTFPLG rPLYADFSHD M. thermophila PAQGVGF-NE LIARLTHSPV QDHTSTNHTL DSNPATFPLN ATLYADFSHD Consensus PAQGVGFANE LIARLTRSPV QDHTSTNHTL DSNPATFPLN ATLYADFSHD Consensus phytase 351 400 SNLVSIFWAL GLYNGTAPLS QTSVESVSQT DGYAAAWTVP FAARAYVEMM A. terreus 9A-1 SNLVSIFWAL GLYNGTKPLS QTTVEDITET DGYAAAWTVP FAARAYIEMM A. terreus cbs A. niger var. awamori ngiisilfal glyngtkpls tttvenitot dgfssawtvp fasrlyvemm A. niger T213 NGIISILFAL GLYNGTKPLS TTTVENITQT DGFSSAWTVP FASRLYVEMM NGIISILFAL GLYNGTKPLS TTTVENITQT DGFSSAWTVP FASRLYVEMM A. niger NRRL3135 A. fumigatus 13073 NSMVSIFFAL GLYNGTEPLS TTSVESAKEL DGYSASWVVP FGARAYFELM A. fumigatus 32722 NSMVSIFFAL GLYNGTGPLS rTSVESAKEL DGYSASWVVP FGARAYFELM NSMVSIFFAL GLYNGTEPLS TTSVESAKEL DGYSASWVVP FGARAYFELM A. fumigatus 58128
A. fumigatus 26906
A. fumigatus 32239 NSMVSIFFAL GLYNGTEPLS rTSVESaKEl DGYSASWVVP FGARAYFELM NGMIPIFFAM GLYNGTEPLS QTSeESTKES NGYSASWAVP FGARAYFELM NSMISIFFAM GLYNGTQPLS mDSVESIQEm DGYAASWTVP FGARAYFELM E. nidulans NTMTSIFaal GLYNGTAKLS TTEIKSIEET DGYSAAWTVP FGGRAYIEMM T. thermophilus NDMMGVLGAL GaYDGVPPLD KTATTDPEEL GGYAASWAVP FAARIYVEKM M. thermophila NSMISIFFAL GLYNGTAPLS TTSVESIEET DGYAASWTVP FGARAYVEMM Consensus NSMISIFFAL GLYNGTAPLS TTSVESIEET DGYSASWTVF FGARAYVEMM Consensus phytase 401 450 QC...... RAEKE PLVRVLVNDR VMPLHGCPTD KLGRCKrDAF A. terreus 9A-1 QC....... RAEKQ PLVRVLVNDR VMPLHGCAVD NLGRCKrDDF A. terreus cbs A. niger T213 QC......QAEQE PLVRVLVNDR VVPLHGCPID aLGRCTrDSF
A. niger NRRL3135 QC.....QAEQE PLVRVLVNDR VVPLHGCPID aLGRCTrDSF A. fumigatus 13073 QC...... KSEKE PLVRALINDR VVPLHGCDVD KLGRCKLNDF QC...... KSEKE PLVRALINDR VVPLHGCDVD KLGRCKLNDF A. fumigatus 32722 A. fumigatus 58128 QC.....KSEKE SLVRALINDR VVPLHGCDVD KLGRCKLNDF A. fumigatus 26906 QC.....KSEKE PLVRALINDR VVPLHGCDVD KLGRCKLNDF QC...... KSEKE PLVRALINDR VVPLHGCAVD KLGRCKLKDF A. fumigatus 32239 QC..... KKE PLVRVLVNDR VVPLHGCAVD KFGRCTLDDW E. nidulans QC......DDSDE PVVRVLVNDR VVPLHGCEVD SLGRCKrDDF T. thermophilus RCsgggggg ggegrQEKDE eMVRVLVNDR VMTLkGCGAD ErGMCTLErF M. thermophila QC----QAEKE PLVRVLVNDR VVPLHGCAVD KLGRCKLDDF Consensus QC..... QAEKE PLVRVLVNDR VVPLHGCAVD KLGRCKRDDF Consensus phytase 451 471 A. terreus 9A-1 VAGLSFAQAG GNWADCF--- -VEGLSFARAG GNWAECF--- -A. terreus cbs A. niger var. awamori VrGLSFARSG GDWAECSA-- -A. niger T213 A. niger NRRL3135 VIGLSFARSG GDWAECFA-- -VrgLsfarsg gdWaecfa-- -A. fumigatus 13073 VKGLSWARSG GNWGECFS-- -A. fumigatus 32722 A. fumigatus 58128 VKGLSWARSG GNWGECFS-- -VKGLSWARSG GNWGECFS-- -A. fumigatus 26906 VKGLSWARSG GNWGECFS-- -

VKGLSWARSG GNSEQSFS-- -VEGLNFARSG GNWKTCFT1- -

T. thermophilus M. thermophila

VrGLSFARQG GNWEGCYAas e IESMAFARGN GKWDlCFA-- -

Consensus Consensus phytase VEGLSFARSG GNWAECFA-- - VEGLSFARSG GNWAECFA...

	1		50
P. involutus (phy		PeseOrn WSPYSPYFPI	AeYkAPPAGC QInQVNIIQR
P. involutus (phy	A2) SVP PniAPK FSI	PeseOrn WSPYSPYFPI	Aeykappagc EinQVNIIQR
	hipirdTSAc LdV	TrDvOgs WSmYSPYFPa	Atyvappasc QinQVHIIQR
T. pubescens	Convostivo nfF	Project WAAYTPYYPV	qaYtPPPkDC KItQVNIIQR
A. pediades	SHOFEFUARO T.PT	PaOntsn WGPYdPFFPV	EpyaAPPEGC tVtQVNLIQR
P. lycii	Storatowne Dir		
Basidio	S-P-R-TAAO LPI	P-O-O WSPYSPYFPV	A-Y-APPAGC QI-QVNIIQR
Basicio			•
	51		100
P. involutus (phy	A1) HGARFPTSGA TTR	IKAGLTK LQGvqnfTD	KFNFIKSfKY dLGnsDLVPF
P. involutus (phy	A2) HGARFPTSGA ATR	IKAGLSK LOSvqnfTDI	KEDEIKZITA GTGEBDTASE
T. pubescens	HGARFPTSGA AKR	IOTAVAK LKAAsnyTDI	llAFVtNyTY sLGqDsLveL
A. pediades	HGARFPTSGA GTR	IOAAVKK LOSAKTYTDI	RLDFLtNyTY tLGhDDLVPF
P. lycii	HGARWPTSGA rSR	QVAAVAK IQMATPETDI	KYEFLNDÍVY KFGVADLLPF
_			
Basidio	HGARFPTSGA ATR	IQAAVAK LQSATDI	KLDFL-N-TY -LG-DDLVPF
	101		150
P. involutus (phy	A1) GAAQSfDAGQ EAF	'ARYSKLV SKNNLPFIRI	dgsdrvvdsa TnwTagfAsA
P. involutus (phy	A2) GAaOSfDAGl EvF	'ARYSKLV SaDNLPFIR	GGSDRVVDTA TNWTAGFASA
T. pubescens	GALOSSEAGO EAF	TRYSSLV SaDELPFVR	SGSDRVVATA INWIAGRALA
A. pediades	GAlQSSQAGE ETF	qRYSfLV SKENLPFVR	A SSSNRVVDSA TNWTEGFSAA
P. lycii	GAnQShQTGt DmY	TRYStLf egGDVPFVR	A AGDQRVVDSS TNWTAGFGDA
_			
Basidio	GA-QSSQAGQ EAF	TRYS-LV S-DNLPFVR	A SGSDRVVDSA TNWTAGFA-A
		•	200
	151		
P. involutus (ph	yA1) ShNTvqPkLn LII	LPQEGNDT LEDNMCPAA	DSDPQvNaWL AVafPSITAR
	yA2) STNAiqPkLd LII	LPQEGNDT LEDNMCPAA	G ESDPQvDaWL AsafPSVTAQ
T. pubescens	SeNSitPvLs VII	SEAGNDT LDDNMCPAA	G DSDPQvNqWL AQFAPPMTAR
A. pediades	ShHvlnPiLf VII	LSESINDT LDDaMCPnA	G SSDPQtGiWt SIYGTPIANR
P. lycii	SgETvlPtLq VVI	LqEeGNcT LcNNMCPnE	v DGDest.tWL GVFAPnITAR
Basidio	s-NTP-L- VII	LSE-GNDT LDDNMCP-A	G DSDPQ-N-WL AVFAPPITAR
			250
	201		-
P. involutus (ph	yai) LNAAAPSVNL TD	DAINLYS LCAFITYSK	E kkSdFCtLFE giPGsFeAFa
P. involutus (ph	yA2) LNAAAPGANL TD:	aDAÍNLVS LCPFMTVSK	E qkSdFCtLFE giPGsFeAFa
T. pubescens	LNAGAPGANL TD	EDTYNLIE LCPFETVAE	E TISEFCDIYE elQAE.dAFa
A. pediades	LNqqAPGANI TA	aDvsNLip LCAFETIVE	E tpSpFCNLFtPEEFaqFe
P. lycii	LNAAAPSANL SD	sDAltLmd MCPFDTLSs	G naSpFCDLFtAEEYvSYe
			m .c.pcnter PERF-AF-
Basidio	LNAAAPGANL TD	-DA-NL LCPFETVS-	ES-FCDLFEPEEF-AF-
	251		300
	251		= :
P. involutus (ph	yAl) YgGDLDKFYG TG	AGGETGEA GGAGIANET	I ARLTHSAVRD NTQTNRTLDA
P. involutus (ph	yaz) YagdldkfYG TG	AGGYTGAA GGAGATNET	L ARLTESAVED HTOTHETLES
T. pubescens		AGGATGAA GGAGATMET	I ARLTaQnVsD HTQTNsTLDS L ARLTemPVRD NTQTNRTLDS
A. pediades	YIGDLDKFYG TG	AGG B T G B A G A G A T M E I	IN WENTERIE AUD MIGINATING
D 2			בי זיים מתוחים ביים או מיים או
P. lycii	YYYDLDKYYG TG	pgnalgpv qgvgyvnei	L ARLTGQAVRD ETQTNRTLDS
Basidio	Yyydldxyyg Tg	pgnalgpv Qgvgyvnei	L ARLT-QAVRD ETQTNRTLDS

350

PCT/DK99/00154

P.	involutus	(phyAl)	SPVTFPLNKT	FYADFSHDNI	MVAVFSAMGL	FIQPAPLSTS	VPNPWRTWIT
D.	involutus	(phyA2)	APCTFPLNKT	MYADFSHDNl	MVAVFSAMGL	FrQSAPLsTS	t PDPNRTWLT
T	nubescens	_	SPETFPLNRT	LYADFSHDNO	MVAIFSAMGL	FNQSAPLDPT	tPDPaRTFLv
Δ.	nediades		SPITFPLDRS	IYADLSHDNO	${\tt MIAIFSAMGL}$	FNQSSPLDPS	f PNPKRTWVT
	lycii		dPaTFPLNRT	FYADFSHDNt	MVPIFAALGL	FNaTA.LDPl	kPDeNRlWVd
Bas	sidio		SP-TFPLNRT	FYADFSHDNQ	MVAIFSAMGL	fnqsapldps	- PDPNRTWVT
			351				400
P.	involutus	(phyA1)	SsLVPFSGRM	VVERLsCf	GT	tkV	RVLVQDqVQP
P.	involutus	(phyA2)	SSVVPFSARM	aVERLsCa	GT	tkV	RVLVQDqVQP
							DT T 122773 110
Α.	pediades		SRLEPFSARM	VtERLlCgrd	GAGTgsggpsri	mrngnvqtfV	RILVNDALQP
P.	lycii		SKLVPFSGHM	tVEKLaC		sgkeaV	RVLVNDAVQP
Bas	sidio		SKLVPFSARM	VVERL-C	GT	v	RVLVNDAVQP
			401			4.4	11
D.	involutus	(phyA1)	LEFCGGDYNG	lCTLAKFVES	QtFARsDGaG	DFEKCFATSa	-
D.	involutus	(phyA2)	LEFCGGDaDG	1 CALDKFVES	QaYARsGGaG	DFEKCLATTV	~
77	pubeccers		TAFCGADESG	VCTLDAFVES	OaYARNDGEG	DFEKCFAT~~	-
Δ.	pediades		LKFCGGDmDS	1CTLEAFVES	QKYAREDGQG	DFEKCFD	-
	lycii		LEFCGG.vDG	vCeLsAFVES	QtyarengQG	DFAKCgfvPs	e ·
			T.PPCCCD_DC	~ CTT.D A FURS	O-VAREDGOG	DFEKCFATP-	-

T. lanuginosa

M. thermophila

Basidio

11/32

```
50
                     KhsdCNSVDh GYQCfPELSH kWGlYAPYFS LqDESPFPlD VPeDCEITFV
                     NhsdCtSVDr GYQCfPELSH kWGlYAPYFS LqDESPFPlD VPdDCHITFV
A. terreus 9al
A. niger var. awamori NqsTCDTVDq GYQCfSEtSH LWGQYAPFFS LANESAISPD VPaGCEVTFa
                    NqsSCDTVDq GYQCfSEtSH LWGQYAPFFS LANESVISPE VPaGCTVTFa
                     GSKSCDTVD1 GYQCsPAtSH LWGQYSPFFS LEDE1SVSSK LPKDCFITLV
A. niger NRRL3135
                     GSKSCDTVDl GYQCsPAtSH LWGQYSPFFS LEDELSVSSK LPkDCRITLV
A. fumigatus 13073
                     GSkSCDTVD1 GYQCsPAtSH LWGQYSPFFS LEDE1SVSSK LPkDCRITLV
A. fumigatus 32722
                     GSkSCDTVDl GYQCsPAtSH LWGQYSPFFS LEDELSVSSK LPkDCRITLV
A. fumigatus 58128
                     GSkACDTVEL GYQCsPGtSH LWGQYSPFFS LEDELSVSSD LPkDCRVTFV
A. fumigatus 26906
A. fumigatus 32239
                      QNHSCNTaDG GYQCfPNVSH VWGQYSPYFS IEQESAISeD VPhGCeVTFV
                      DSHSCNTVEG GYQCrPEISH SWGQYSPFFS LADQSEISPD VPQNCKITFV
E. nidulans
T. thermophilus
                      ----nvDIAR hwgQYSPFFS LAEVSEISPA VPkGCRVeFV
                      ESRPCDTpDl GFQCgTAISH FWGQYSPYFS VPsElDaS.. IPdDCeVTFa
T. lanuginosa
                      xSxPxrxtAA qLPipxQxqx xWSPYSPYFP VAxyxA.... pPaGCQIxqV
M. thermophila
Basidio
           Consensus NSHSCDTVDG GYQC-PEISH LWGQYSPFFS LADESAISPD VP-GCRVTFV
                Fcp10 NSHSCDTVDG GYQCFPEISH LWGQYSPFFS LADESAISPD VPKGCRVTFV
                       QVLARHGARS PThSKTKAYA AtlaAlQKSA TaFpGKYAFL QSYNYSLDSE
                       QVLARHGARS PTdSKTKaYA AtlaAlQKNA TalpGKYAFL KSYNYSMGSE
 A. terreus 9al
 A. niger var. awamori QVLSRHGARY PTESKGKKYS ALIEEIQQNv TtFDGKYAFL KTYNYSLGAD
                     QVLSRHGARY PTdSKGKKYS ALIEEIQQNA TtFDGKYAFL KTYNYSLGAD
 A. niger NRRL3135
                       QVLSRHGARY PTSSKSKKYK KLVtAIQANA TdFKGKFAFL KTYNYTLGAD
 A. fumigatus 13073
                       QVLSRHGARY PTSSKSKKYK KLVLAIQANA TdFKGKFAFL KTYMYTLGAD
                       QVLSRHGARY PTSSKSKKYk kLVtAIQANA TdFKGKFAFL KTYNYTLGAD
 A. fumigatus 32722
 A. fumigatus 58128
                       QVLSRHGARY PTSSKSKKYk kLVtAIQaNA TdFKGKFAFL KTYNYTLGAD
                       QVLSRHGARY PTASKSKKYK KLVTAIQKNA TEFKGKFAFL ETYNYTLGAD
 A. fumigatus 26906
                       QVLSRHGARY PTESKSKAYS GLIEAIQKNA TSFWGQYAFL ESYNYTLGAD
 A. fumigatus 32239
                       QLLSRHGARY PTSSKTELYS qLIsrIQKtA TaYKGYYAFL KdYrYqLGAN
 E. nidulans
                       QVLSRHGARY PTAhKSEVYA ELLQTIQDtA TeFKGDFAFL RdYaYhLGAD
 T. thermophilus
                       QVLSRHGARa PTlkRAasYv DLIdriHhGA isYgPgYEFL RTYDYTLGAD
 T. lanuginosa
                       NIIQRHGARF PTSGaAtRiq AaVakLQsax xxtDPKLDFL xnxtYxLGxD
 M. thermophila
 Basidio
            Consensus QVLSRHGARY PTSSKSKKYS ALI-AIQKNA T-FKGKYAFL KTYNYTLGAD
                Fcplo QVLSRHGARY PTSSKSKKYS ALIEAIQKNA TAFKGKYAFL KTYNYTLGAD
                        ELTPFGrNQL rDlGaQFYeR YNAL.TRhin PFVRATDASR VhESAEKFVE
                        101
                       NLTPFGrNQL qDlGaQFYRR YDTL.TRhIn PFVRAADSSR VhESAEKFVE
  A. terreus 9al
  A. terreus cbs
  A. niger var. awamori DLTPFGEQEL VNSGIKFYQR YESL.TRnII PFIRSSGSSR VIASGEKFIE
                      DLTPFGEQEL VNSGIKFYQR YESL.TRnIV PFIRSSGSSR VIASGKKFIE
  A. niger NRRL3135
                       DLTPFGEQQL VNSGIKFYQR YKAL.ARSVV PFIRASGSDR VIASGEKFIE
  A. fumigatus 13073
                     DLTPFGEQQL VNSGIKFYQR YKAL.ARSVV PFIRASGSDR VIASGEKFIE
                     DLTPFGEQQL VNSGIKFYQR YKAL.ARsVV PFIRASGSDR VIASGEKFIE
  A. fumigatus 32722
  A. fumigatus 58128
                        DLTAFGEQQL VNSGIKFYQR YKAL.ARSVV PFIRASGSDR VIASGEKFIE
  A. fumigatus 26906
                        DLTPFGEQQM VNSGIKFYQK YKAL.AgsVV PFIRSSGSDR VIASGEKFIE
  A. fumigatus 32239
                        DLTiFGENQM VDSGaKFYRR YKnL.ARknt PFIRASGSDR VVASAEKFIN
  E. nidulans
                        DLTPFGENQM IQlGIKFYDH YKSL.ARDAV PFVRCSGSDR VIASGrlFIE
  T. thermophilus
                        NLTRFGEEQM MESGrQFYHR YREq.AReIV PFVRAAGSAR VIASAEfFnr
```

ELTREGOOOM VNSGIKFYRR YRAL.ARKSI PFVRTAGODR VVhSAENFEQ

DLvPFGAxQs sQAGqEaFtR YsxLvSxdnL PFVRASGSDR VVDSAtNWtA

Consensus DLTPFGEQQM VNSGIKFYRR YKAL-AR-IV PFVRASGSDR VIASAEKFIE

12/32
Fcp10 DLTPFGEQQM VNSGIKFYRR YKAL.ARKIV PFVRASGSDR VIASAEKFIE

```
GFQTARqDDh hAnphQPSPr VDVaIPEGsA YNNTLEHSLC TAFEs...St
A. terreus 9al
                      GFQNARQGDP hanphQPSPr VDVVIPEGtA YNNTLEHSIC TAFEa...St
A. terreus cbs
A. niger var. awamori GFQSTKLkDP rAqpgQSSPk IDVVISEASS SNNTLDpGtC TvFEd...SE
                     GFQSTKLkDP rAqppgQSSPk IDVVISEASS aNNTLDpGtC TvFed...SE
A. niger NRRL3135
                     GFQQAKLADP gAt.nRAAPa ISVIIPESeT FNNTLDHGVC TkFEa...SQ
A. fumigatus 13073
                     GFQQAKLADP gAt.nRAAPa ISVIIPESeT FNNTLDHGVC TKFEa...SQ
A. fumigatus 32722
                     GFQqAKLADP gAt.nRAAPa ISVIIPESeT FNNTLDHGVC TKFEa...SQ
A. fumigatus 58128
                      GFQQAKLADP gAt.nRAAPa ISVIIPESeT FNNTLDHGVC TkFEa...SQ
A. fumigatus 26906
                      GFQqANVADP gAt.nRAAPV ISVIIPESET YNNTLDHSVC TnFEa...SE
A. fumigatus 32239
                      GFRkAQLhDh g.s.gQATPV VNVIIPEidG FNNTLDHStC vSFEn...dE
E. nidulans
                      GFQSAKVlDP hSdkhDAPPt INVIIeEGpS YNNTLDtGsC PvFEd...Ss
T. thermophilus
                      GFQdAKdrDP rsnkdQAePV INVIISEEtG sNNTLDgltC PAaEe...Ap
T. lanuginosa
                      GFHSALLADR gStvrPTlPy dmVVIPETaG aNNTLHNDLC TAFEegPySt
M. thermophila
                      GFaxA..... ..sxntxxPx LxVILSExg. .NDTLDDNMC .....PxAG
Basidio
           Consensus GFQSAKLADP -A---QASPV INVIIPEG-G YNNTLDHGLC TAFE--P-SE
               Pcp10 GFQSAKLADP GANPHQASPV INVIIPEGAG YMNTLDHGLC TAFEE...SE
                      201
                      VGDDavANFT AVFAPAIaqR LEAGLPGVQL StDDVVNLMA MCPFETVSlT
A. terreus 9al
                      VGDAaADNFT AVFAPAIakR LEAGLPGVQL SADDVVNLMA MCPFETVSlT
A. terreus cbs
A. niger var. awamori LADtVEANFT AtFAPSIRQR LEndLSGVtL TDtEVtyLMD MCSFDTIStS
A. niger NRRL3135 LADTVEANFT ATFVPSIRGR LENGLSGVTL TDTEVTYLMD MCSFDTISTS
                    LGDEVAANFT ALFAPdIRAR aEkhLPGVtL TDEDVVSLMD MCSFDTVArT
A. fumigatus 13073
A. fumigatus 32722 LGDEVAANFT ALFAPdIRAR aEkhLPGVtL TDEDVVSLMD MCSFDTVArT
A. fumigatus 58128 LGDEVAANFT ALFAPdIRAR aEkhLPGVtL TDEDVVSLMD MCSFDTVArT
A. fumigatus 26906 LGDEVAANFT ALFAPdIRAR aKkhLPGVtL TDEDVVSLMD MCSFDTVArT
                     LGDEVEANFT ALFAPAIRAR IEKhLPGVQL TDDDVVSLMD MCSFDTVArT
A. fumigatus 32239
                      radeleanft aimgppirkr lendlpgikl tnenvlylmd mcsfdtmart
E. nidulans
                      gGHDaQEKFA kqFAPAIlEK IKDhLPGVDL AvsDVpyLMD LCPFETLArn
T. thermophilus
                      .DptqpAEFl qVFGPRVlkK ItkhMPGVNL TlEDVplFMD LCPFDTVGsd
T. lanuginosa
                      IGDDaQDtYl StFAGPItAR VNAnLPGaNL TDADtVaLMD LCPFETVAsS
M. thermophila
                      dSDpqxnxWl AVFAPPItAR LNAaaPGaNL TDxDaxNLxx LCPFETVS..
Basidio
           Consensus LGDDVEANFT AVFAPPIRAR LEA-LPGVNL TDEDVVNLMD MCPFDTVA-T
               Fcp10 LGDDVEANFT AVFAPPIRAR LEAHLPGVNL TDEDVVNLMD MCPFDTVART
                      dD...Aht... ......LSPF CDLFTa..tE WtQYNYLlSL dKYYGYGGGN
 A. terreus 9al
                      dD..Aht... .....LSPF CDLFTa..aE WtQYNYLlSL dKYYGYGGGN
 A. terreus cbs
 A. niger var. awamori Tv..DTK... .....LSPF CDLFTH..dE WiHYDYLQSL kKYYGHGAGN
 A. niger NRRL3135 Tv..DTK... .....LSPF CDLFTH..dE WinyDyLQSL kKYYGHGAGN
                     SD..ASQ... .....LSPF CQLFTH..nE WKKYNYLQSL GKYYGYGAGN
 A. fumigatus 13073
                     SD..ASQ... .....LSPF CQLFTH..nE WKKYNYLQSL GKYYGYGAGN
 A. fumigatus 32722
                      SD..ASQ.....LSPF CQLFTH..nE WKKYNYLQSL GKYYGYGAGN
 A. fumigatus 58128
                      SD..ASQ... LSPF CQLFTH..nE WKKYNYLQSL GKYYGYGAGN
 A. fumigatus 26906
                      AD..ASE... .....LSPF CAIFTH..nE WKKYDYLQSL GKYYGYGAGN
 A. fumigatus 32239
                      AH..GTE... LSPF CAIFTE..kE WlQYDYLQSL sKYYGYGAGS
 E. nidulans
                      ht..DT.... LSPF CALSTQ..eE WqaYDYYQSL gKYYGnGGGN
 T. thermophilus
                      PvlfPrQ.....LSPF CHLFTa..dD WmaYDYYYTL dKYYSHGGGS
 T. lanuginosa
                      SsdpATadag ggngrpLSPF CrLFSE..sE WraYDYLQSV gKWYGYGPGN
 M. thermophila
                       ...... ...xexxSxF CDLFexxpeE FxaFxYxgdL dKFYGtGyGQ
 Basidio
            Consensus SD--ATQ--- ----LSPF CDLFTH---E W-QYDYLQSL -KYYGYGAGN
                Fcp10 SD..ATQ... .....LSPF CDLFTH..DE WIQYDYLQSL GKYYGYGAGN
                                                                         350
                       PLGPvQGVGW anelmarltr A.PVHDHTCv nntldaspat fplnatlyad
 A. terreus 9al
                       PLGPvQGVGW aneliarltr s.pvHDHTCv nntlDanpat fpLnatlyaD
 A. terreus cbs
 A. niger var. awamori PLGPTQGVGY aNELIARLTH S.PVHDDTSS NHTLDSNPAT FPLNSTLYAD
```

A. niger NRRL3135

PLGPTQGVGY aNELIARLTH S.PVHDDTSS NHTLDSSPAT FPLNSTLYAD

A. fumigatus 13073

13/32

PLGPAQGIGF tNELIARLTR S.PVQDHTST NSTLvSNPAT FPLNATMYvD

```
A. fumigatus 32722
                     PLGPAQGIGF tNELIARLTR S.PVQDHTST NSTLvSNPAT FPLNATMYvD
A. fumigatus 58128
                     PLGPAQGIGF tHELIARLTR S.PVQDHTST NSTLVSNPAT FPLNATMYvD
A. fumigatus 26906
                     PLGPAQGIGF tHELIARLTR S.PVQDHTST NSTLvSNPAT FPLNATMYvD
A. fumigatus 32239
                     PLGPAQGIGF tNELIARLTN S.PVQDHTST NSTLDSDPAT FPLNATIYVD
                     PLGPAQGIGF tNELIARLTQ S.PVQDNTST NHTLDSNPAT FPLDrkLYAD
E. nidulans
                     PLGPAQGVGF VNELIARMTH S.PVQDYTTV NHTLDSNPAT FPLNATLYAD
T. thermophilus
T. lanuginosa
                     AFGPSRGVGF VNELIARMTG N1PVKDHTTV NHTLDdNPET FPLDAVLYAD
M. thermophila
                     PLGPTQGVGF vNELLARLA. GVPVRDgTST NRTLDGDPrT FPLGrPLYAD
Basidio
                     PLGPvQGVGY iNELLARLTx qa.VRDNTqT NRTLDSSPxT FPLNrTFYAD
           Consensus PLGPAQGVGF -NELIARLTH S-PVQDHTST NHTLDSNPAT FPLNATLYAD
              Fcp10 PLGPAQGVGF VNELIARLTH S.PVQDHTST NHTLDSNPAT FPLNATLYAD
                     FSHDSnLVSI FWALGLYNGT aPLSqTSVE. .SvsQTDGYA AAWTVPFAAR
A. terreus 9al
                     FSHDSnLVSI FWALGLYNGT kPLSqTTVE. .ditrTDGYA AAWTVPFAAR
A. terreus cbs
A. niger var. awamori FSHDNGIISI LFALGLYNGT kPLSTTTVE. .NitQTDGFS SAWTVPFASR
A. niger NRRL3135 FSHDNGIISI LFALGLYNGT kPLSTTTVE. .NitQTDGFS SAWTVPFASR
                   FSHDNSMVSI FFALGLYNGT ePLSTTSVE. .SaKElDGYS ASWVVPFGAR
A. fumigatus 13073
A. fumigatus 32722
                   FSHDNSMVSI FFALGLYNGT qPLSrTSVE. .SaKElDGYS ASWvVPFGAR
A. fumigatus 58128
                   FSHDNSMVSI FFALGLYNGT ePLSTTSVE. .SaKElDGYS ASWvVPFGAR
A. fumigatus 26906
                   FSHDNSMVSI FFALGLYNGT ePLSrTSVE. .SaKElDGYS ASWvVPFGAR
                   FSHDNGMIPI FFAMGLYNGT ePLSqTSeE. .StKESNGYS ASWAVPFGAR
A. fumigatus 32239
                     FSHDNSMISI FFAMGLYNGT qPLSmdSVE. .SiQEmDGYA ASWTVPFGAR
E. nidulans
                     FSHDNTMtSI FaALGLYNGT akLSTTeIK. .SiEETDGYS AAWTVPFGGR
T. thermophilus
                     FSHDNTMtGI FSAMGLYNGT kPLSTSkIQP pTgAAADGYA ASWTVPFAAR
T. lanuginosa
                     FSHDNdMMGV LgALGaYDGv pPLdkTA..R rdpEElGGYA ASWAVPFAAR
M. thermophila
Basidio
                     FSHDNgMVAI FSAMGLFNgS aPLdPSxpDP nrt....Wv TSklVPFSAR
          Consensus FSHDNTMVSI FFALGLYNGT -PLSTTSVEP -S-EETDGYA ASWTVPFAAR
              Fcp10 FSHDNTMVSI FFALGLYNGT KPLSTTSVE. .SIEETDGYA ASWTVPFAAR
                     401
                     AYVEMMQC.. ra..... .....EKEPL VRVLVNDRVM PLHGCPtDKL
A. terreus 9al
                     AYIEMMQC.. ra..... EKQPL VRVLVNDRVM PLHGCAVDNL
A. terreus cbs
A. niger var. awamori lyvemmQC.. Qa...... EQEPL VRVLVNDRVV PLHGCPIDaL
                     lyvemmqc.. Qa..... .....EQEPL VRVLVNDRVV PLHGCPVDaL
A. niger NRRL3135
                     AYFETMQC.. Ks..... ....EKEPL VRaLINDRVV PLHGCDVDKL
A. fumigatus 13073
                     AYFETMQC.. Ks..... EKEPL VRaLINDRVV PLHGCDVDKL
A. fumigatus 32722
                     AYFETMQC.. Ks..... .....EKESL VRALINDRVV PLHGCDVDKL
A. fumigatus 58128
                     AYFETMQC.. Ks..... EKEPL VRaLINDRVV PLHGCDVDKL
A. fumigatus 26906
                     AYFETMQC.. Ks..... EKEPL VRaLINDRVV PLHGCAVDKL
A. fumigatus 32239
                     AYFELMQC.. E...... KKEPL VRVLVNDRVV PLHGCAVDKF
E. nidulans
                     AYIEMMQC.. Dd...... SDEPV VRVLVNDRVV PLHGCEVDsL
T. thermophilus
                     AYVELLRC.. Etetsseeee EG...EDEPF VRVLVNDRVV PLHGCrVDRW
T. lanuginosa
                     iYVEkMRC.. sgggggggg EGrqeKDEeM VRVLVNDRVM TLkGCGaDEr
M. thermophila
                     mvVErLxCxx xgtxxxxxxx xxxxxxxxx VRVLVNDaVq PLEfCGgDxd
Basidio
           Consensus AYVEMMOC-- E----- EG---EKEPL VRVLVNDRVV PLHGCGVDKL
              Fcp10 AYVEMMQC.. EA...... EKEPL VRVLVNDRVV PLHGCGVDKL
```

	•	451			82
a	terreus 9al	GRCKrDAFVA	GLSFAQAG	GNWADCF	
	terreus cbs	GRCKrDDFVE		GNWAECF	
	niger var. awamori			GDWAECsA	
	niger NRRL3135	GRCtrDsFVr		GDWAECFA	
	fumigatus 13073	GRCKINDFVK		GNWGECFS	
	fumigatus 32722	GRCKINDFVK		GNWGECFS	
	fumigatus 58128	GRCKINDFVK		GNWGECFS~~	
	fumigatus 26906		GLSWARSG	GNWGECFS~~	
	fumigatus 32239		GLSWARSG	GNSEQSFS~~	
	nidulans		GLNFARSG	GNWKtCFT1~	
	thermophilus	GRCKrDDFVr	GLSFARqG	GNWEGCYAas	
	lanuginosa	GRCRrDEWIK	GLTFARqG	GHWDrCF	
м.	thermophila	GmCtlErFIE	SMAFARGN	GKWDlCFA~~	
	sidio	GxCtlDAFVE	SqxYAReDgq	GDFEKCFAtp	XX
	Consensus	GRCK-DDFVE	GLSFARSG	GNWEECFA	
	Fcp10		GLSFARSG		

	1	mm!	-MCDVCDVED	T 7 E2267	50
P. involutus (phyAl)	~~~~~~	~Fripesedk	-MCDACDAED	LAEykA LAEykA	praccoingv
P. involutus (phyA2)		~Fsipesedk	IMPETPETE	aAtyvA	nPascolingV
T. pubescens		~TDACKDAGO	CVVQTV CKWA	VqAyTP	PERDCATURA
A. pediades		~prrpPQrqo	PHYGDAGEED	VEpyAA	PERCHUTCU
P. lycii	** -3@*****	~LPIPAUNIS	TIMGETOFFF	LqDESPFPlD	DESCRIPTION
A. terreus 9al	KnsdCNSVDn	CYCCEPELSA	LWG1 VADVES	LqDESPFP1D	VPDCHITEV
A. terreus cbs	NASCESVDE	GIÓCIATRES	TWCOVAPEE	LANESAISPD	VPacceurra
A. niger var. awamori	Masichiana	CAUCTREFER	I.WGOVAPEES	LANESVISPD	VPacCRVTFa
A. niger T213	Mascolomba	CAUCESECRE	LWCOVAPFES	LANESVISPE	VPaGCRVTFa
A. niger NRRL3135 A. fumigatus ATCC13073	udeschiand	GYOCEDAESH	LWGOVSPFFS	LEDELSVSSK	LPKDCRITLV
A. fumigatus ATCC32722	GSYSCDIADI	GYOCEPALSH	LWGOYSPEES	LEDELSVSSK	LPKOCRITLV
A. fumigatus ATCC58128	GSKSCDIVDI	GYOCEPALSH	LWGOYSPEES	LEDELSVSSK	LPKDCRITLV
A. fumigatus ATCC26906	GSKSCDIVDI	GYOCEPATSH	LWGOYSPFFS	LEDELSVSSK	LPKDCRITLV
A. fumigatus ATCC32239	GSKACDTVEI	GYOCEPGESH	LWGOYSPFFS	LEDELSVSSD	LPKDCRVTFV
E. nidulans	ONUSCATADO	GYOCEPNVSH	VWGOYSPYFS	IEQESAISeD	VPhGCeVTFV
T. thermophilus	DSHSCNTVEG	GYOCYPETSH	sWGOYSPFFS	LADQSEISPD	VPONCKITFV
T. lanuginosa		~~~nvDTAR	hWGOYSPFFS	LAEVSEISPA	VPKGCRVeFV
M. thermophila	ESSPCDToD1	GFOCGTAISH	FWGOYSPYFS	VPsElDaS	IPDDCeVTFa
M. Cheimophila	20.11 021 221	0. 402			
Consensus Seq. 11	NSHSCDTVD-	GYOC-PEISH	LWGQYSPFFS	LADESAISPD	VPKGCRVTFV
COMBERGED DOQ. EL	,		-		
	51				100
P. involutus (phyA1)	NIIGRHGARF	PTSGaTtRik	AgLtKLQgvq	nftDAKFnFI	KSFKYdLGns
P. involutus (phyA2)	NIIGRHGARF	PTSGaAtRik	AgLsKLQsvq	nftDPKFDFI	KSFtYdLGTs
T. pubescens	HIIGRHGARF	PTSGaAKRig	TaVAKLKaaS	nytDPlLAFV	tnYtYSLGqD
A. pediades	NIIGRHGARF	PTSGaGtRig	AaVKKLQsak	TytDPRLDFL	tnYtYTLGhD
P. lycii	NLIGRHGARW	PTSGarsRov	AaVAKIQmar	PftDPKYEFL	NdfvYkFGvA
A. terreus 9al	OVLARHGARS	PThSKTKaYA	AtlAalOKSA	TafpGKYAFL	QSYNYSLDSE
A. terreus cbs	OVI.ARHGARS	PTdSKTKaYA	AtlAaIOKNA	TalpGKYAFL	KSYNYSMGSE
A. niger var. awamori	OVLSRHGARY	PTeSKGKKYS	ALIEeIQQNv	TtFDGKYAFL	KTYNYSLGAD
A. niger T213	QVLSRHGARY	PTeSKGKKYS	ALIEeIQQNv	TtFDGKYAFL	KTYNYSLGAD
A. niger NRRL3135	QVLSRHGARY	PTdSKGKKYS	ALIEEIQQNA	TtFDGKYAFL	KTYNYSLGAD
A. fumigatus ATCC13073	QVLSRHGARY	PTSSKSKKYk	kLVtaIQaNA	Tdfkgkfafl	KTYNYTLGAD
A. fumigatus ATCC32722	QVLSRHGARY	PTSSKSKKYk	kLVtaIQaNA	Tdfkgkfafl	KTYNYTLGAD
A. fumigatus ATCC58128	QVLSRHGARY	PTSSKSKKYk	kLVtaIQaNA	Tdfkgkfafl	KTYNYTLGAD
A. fumigatus ATCC26906	QVLSRHGARY	PTSSKSKKYk	kLVtaIQaNA	TdfKGKFAFL	KTINYTLGAD
A. fumigatus ATCC32239	QVLSRHGARY	PTASKSKKYK	kLVtaIQKNA	TeFKGKFAFL	ETINYTLGAD
E. nidulans	QVLSRHGARY	PTeSKSKaYS	GLIEAIQKNA	TsFwGQYAFL	ESYNYTLGAD
T. thermophilus	QLLSRHGARY	PTSSKTELYS	qLIsRIQKtA	TaYKGyYAFL	Kairiare
T. lanuginosa	QVLSRHGARY	PTAhKSEVYA	ELLQRIQDEA	TeFKGDFAFL	ROYAYALGAD
M. thermophila	QVLSRHGARa	PTlkRAasYv	DLIDRIHHGA	isYgPgYEFL	RYDYTLGAD
•					VENUATION CAD
Consensus Seq. 11	QVLSRHGARY	PTSSKSKKYS	ALIERIOKNA	T-FKGKYAFL	KIINIILIGAD
			•		
101					
150					TEMPORENTIAL D
P. involutus (phyA1)	DLvPFGAaQs	fDAGqEaFaR	YSKLVSKNIL	PFIRAGSDR	AADOVERHEY
P. involutus (phyA2)	DLvPFGAaQs	fDAGLEvFaR	YSKLVSSDDL	PFIRSdGSDR	AADIVENACU
T. pubescens	sLveLGAtQs	sEAGqEaFtR	YSSLVSabeL	PFVRASGSDR	AANIMIMEN
A. pediades	DLvPFGAlQs	sQAGeEtFQR	YsfLvskenL	PFVRASSSNR	VVDSACNWCE
P. lycii	DL1PFGANQs	hQTGtDMYtR	YSTLIEGGOV	PFVRAAGdQR	VANDOCUMCA
A. terreus 9al	ELTPFGINQL	rDlGaQFYeR	YNAL.TRHIN	PFVRATDASR	VIRDAERFVE
A. terreus cbs	NLTPFGrnQL	qD1GaQFYRR	YDTL.TRHIN	PFVRAADSSR	VDESAERFVE
A. niger var. awamori	DLTPFGEQEL	VNSGIKFYOR	YESL.TRNII	PFIRSSGSSR	VIASGERFIE
A. niger T213	DLTPFGEQEL	VNSGIKFYOR	YESL.TRNII	PFIRSSGSSR	VIASGERFIE
A. niger NRRL3135	DLTPFGEQEL	VNSGIKFYQR	YESL.TRNIV	PFIRSSGSSR	VIASGRAPIE
A. fumigatus ATCC13073	DLTPFGEQQL	VNSGIKFYQR	YKAL.ARSVV	PFIRASGSDR	ATVOCEVETE
A. fumigatus ATCC32722	DLTPFGEQQL	VNSGIKFYQR	YKAL.ARSVV	PFIRASGSDR	ATVOCEABLE
A. fumigatus ATCC58128	DLTPFGEQQL	VNSGIKFYQR	YKAL.ARSVV	PFIRASGSDR	* TWOGEVE IE
	•		O A		

WO 99/48380 PCT/DK99/00154

```
A. fumigatus ATCC26906 DLTAFGEQQL VNSGIKFYQR YKAL.ARSVV PFIRASGSDR VIASGEKFIE A. fumigatus ATCC32239 DLTPFGEQQM VNSGIKFYQK YKAL.AGSVV PFIRSSGSDR VIASGEKFIE
                        DLTiFGENOM VDSGaKFYRR YKnL.ARKnt PFIRASGSDR VVASAEKFIN
E. nidulans
                        DLTPFGENOM IQIGIKFYnH YKSL.ARNAV PFVRCSGSDR VIASGRIFIE
T. thermophilus
                        NLTRFGEEQM MESGrQFYHR YREq.AREIV PFVRAAGSAR VIASAEffnr
T. lanuginosa
                        ELTRtGQQQM VNSGIKFYRR YRAL.ARKsI PFVRTAGqDR VVhSAENFtQ
M. thermophila ~
                        DLTPFGENOM VNSGIKFYRR YKAL-ARNIV PFVRASGSDR VIASAEKFIE
Consensus Seq. 11
                                                                            200
                        GFaSA..... ..shNtvqPk LNLILPQ..T gNDTLEDNMC PAaGD.....
P. involutus (phyA1)
                        GFaSA..... ..srNaiqPk LDLILPQ..T gNDTLEDNMC PAaGE.....
P. involutus (phyA2)
                        GFala.... ..ssNsiTPV LSVIISE..A gNDTLDDNMC PAaGD.....
T. pubescens
                        GFsAA..... ..shHvlNPI LfVILSE..S LNDTLDDAMC PnaGs.....
A. pediades
                        GFgdA..... ..sgEtvlPt LQVVLQE..E gNcTLcNNMC PnevD.....
P. lycii
                        GFQTARqDDh hAnpHQPSPr VDVaIPEGSA YNNTLEHSLC TAFEs...ST
A. terreus 9al
                        GFQNARqGDP hAnpHQPSPr VDVVIPEGTA YNNTLEHSIC TAFEA...ST
A. terreus cbs
A. niger var. awamori GFQSTKLkDP rAqpgQSSPk IDVVISEASS SNNTLDpGtC TvFED...Se
                         GFQSTKLkDP rAqpgQSSPk IDVVISEASS sNNTLDpGtC TvFED...Se
A. niger T213
                         GFQSTKLkDP rAqpgQSSPk IDVVISEASS SNNTLDpGtC TvFED...Se
A. niger NRRL3135
A. fumigatus ATCC13073 GFQqAKLADP gAt.NRAAPa ISVIIPESeT FNNTLDHGVC TKFEA...Sq
A. fumigatus ATCC32722 GFQQAKLADP gAt.NRAAPa ISVIIPESeT FNNTLDHGVC TKFEA...Sq
A. fumigatus ATCC58128 GFQQAKLADP gAt.NRAAPa ISVIIPESeT FNNTLDHGVC TKFEA...Sq
A. fumigatus ATCC26906 GFQQAKLADP GAt.NRAAPa ISVIIPESeT FNNTLDHGVC TKFEA...SQ
A. fumigatus ATCC32239 GFQQANVADP GAt.NRAAPV ISVIIPESET YNNTLDHSVC ThFEA...Se
                         GFRkAQLhDh g.s.gQATPV VNVIIPEidG FNNTLDHStC vSFEN...de
                         GFQSAKVlDP hSdkHDAPPt INVIIeEGPS YNNTLDtGsC PvFED...SS
 T. thermophilus
                         GFQdAKdrDP rSnkDQAePV INVIISEETG sNNTLDgltC PAaEE...AP
 T. lanuginosa
                         GFHSALLADR gStvRPTlPy dmVVIPETAG aNNTLHNDLC TAFEEgpyST
 M. thermophila
                         GFQSAKLADP -A--HQASPV INVIIPEGSG YNNTLDHGLC TAFED---ST
 Consensus Seq. 11
                         .SDpqvnaWl AVafPSItAR LNAaaPSVNL TDtDafNLVs LCAFITVSK.
 P. involutus (phyA1)
                         .SDpqvDaWl AsafPSVtAQ LNAaaPGaNL TDADafNLVs LCPFmTVSK.
 P. involutus (phyA2)
                         .SDpqvnQwl AqFAPPMtAR LNAgaPGaNL TDtDtyNLLt LCPFETVAt.
 T. pubescens
                         .SDpqtGiWT SIYGTPIanR LNqqaPGaNI TAADVSNLIP LCAFETIVK.
 A. pediades
                         .GDESt.twl GVFAPnItAR LNAaaPSaNL SDsDaLtLMD MCPFDTLSs.
 P. lycii
                         VGDDAVANFT AVFAPAIaqR LEAGLPGVQL StDDVVNLMA MCPFETVSlT
 A. terreus 9al
                         VGDAAADNFT AVFAPAIAKR LEAGLPGVQL SADDVVNLMA MCPFETVSlT
 A. terreus cbs
 A. niger var. awamori LADtvEANFT AtFAPSIRQR LEndLSGVtL TDtEVtyLMD MCSFDTIStS
                        LADTVEANFT ATFAPSIRGR LENGLSGVTL TOTEVTYLMD MCSFDTISTS
 A. niger T213
                         LADIVEANFT ATTVPSIRGR LENGLSGVIL TDTEVTYLMD MCSFDTISTS
 A. niger NRRL3135
 A. fumigatus ATCC13073 LGDEVAANFT ALFAPdIRAR aEKhLPGVtL TDEDVVSLMD MCSFDTVART
 A. fumigatus ATCC32722 LGDEVAANFT ALFAPdIRAR aEKhLPGVtL TDEDVVSLMD MCSFDTVART
 A. fumigatus ATCC58128 LGDEVAANFT ALFAPdIRAR aEkhLPGVtL TDEDVVSLMD MCSFDTVART
 A. fumigatus ATCC26906 LGDEVAANFT ALFAPdIRAR aKkhLPGVtL TDEDVVSLMD MCSFDTVART
 A. fumigatus ATCC32239 LGDEVEANFT ALFAPAIRAR IEKhLPGVQL TDDDVVSLMD MCSFDTVART
                         radeieanft almgppirkr LendLpgikl TNENVIYLMD MCSFDTMART
 E. nidulans
                         gGHDAQEKFA kqFAPAIlEK IKDhLPGVDL AVsDVpyLMD LCPFETLARn
 T. thermophilus
                         .DptqpAEF1 qVFGPRVlkK ItkhMPGVNL TlEDVplFMD LCPFDTVGsd
  T. lanuginosa
                         IGDDAQDtyl StFAGPItAR VNAnLPGaNL TDADtVaLMD LCPFETVASS
 M. thermophila
                          LGDDAEANFT AVFAPPIRAR LEA-LPGVNL TDEDVVNLMD MCPFDTVART
 Consensus Seq. 11
                          251
                          .....ekkSdF CtLFegiPGs FeaFAYggdL dKFYGtGyGQ
  P. involutus (phyA1)
                          .....eqkSdF CtLFegiPGs FeaFAYagdL dKFYGtGyGQ
  P. involutus (phyA2)
                          ............errSeF CDIYeelqAE .daFAYnadL dKFYGtGyGQ
  T. pubescens
                          .....etpSPF CNLF..TPEE FaQFEYFgdL dXFYGtGyGQ
  A. pediades
                          .....gnaSPF CDLF..TAEE YvsYEYYYdL dKYYGtGPGN
  P. lycii
                          dD..Aht.....LSPF CDLF..TAtE WtQYNYLlSL dKYYGYGGGN
  A. terreus 9al
                          dD..Aht... .....LSPF CDLF..TAAE WtQYNYLlSL dKYYGYGGGN
  A. terreus cbs
                          TV..DTK... .....LSPF CDLF..ThDE WiHYDYLQSL kKYYGHGAGN
  A. niger var. awamori
```

Fig. 8B

```
17/32
                         Tv..DTK.....LSPF CDLF..ThDE WiHYDYLRSL kKYYGHGAGN
A. niger T213
                         Tv..DTK... .....LSFF CDLF..ThDE WiNYDYLQSL kKYYGHGAGN
A. niger NRRL3135
A. fumigatus ATCC13073 SD..ASQ... .....LSFF CQLF..ThNE WKKYNYLQSL gKYYGYGAGN
A. fumigatus ATCC32722 SD..ASQ... .....LSFF CQLF..ThNE WKKYNYLQSL gKYYGYGAGN
A. fumigatus ATCC58128 SD..ASQ... .....LSPF CQLF..ThNE WKKYNYLQSL GKYYGYGAGN
A. fumigatus ATCC26906 SD..ASQ... .....LSPF CQLF..ThNE WKKYNYLQSL GKYYGYGAGN
A. fumigatus ATCC32239 AD..ASE... .....LSPF CAIF..ThNE WKKYDYLQSL gKYYGYGAGN
                     AH..GTE...LSPF CAIF..TEKE WlQYDYLQSL SKYYGYGAGS
ht..DT...LSPF CALS..TGEE WQAYDYYQSL GKYYGNGGGN
PvlfprQ...LSPF CHLF..TADD WmayDYYYTL dKYYSHGGGS
E. nidulans
T. thermophilus
T. lanuginosa
                        SsdpATadag ggngrpLSPF CrLF..SESE WraYDYLQSV gKWYGYGPGN
M. thermophila
                         SD--ATQ--- -----LSPF CDLF--TADE W-QYDYLQSL -KYYGYGAGN
Consensus Seq. 11
                         301
                         eLGPvQGVGY vNELIARLIN S.AVRDNTqT NRTLDASPvT FPLNkTFYAD
P. involutus (phyA1)
                         ALGPVQGVGY INELLARLIN S.AVNDNTQT NRTLDAaPDT FPLNkTMYAD
P. involutus (phyA2)
                        PLGPvQGVGY iNELIARLTa q.nVsDHTqT NsTLDSSPET FPLNrTLYAD
T. pubescens
                        PLGPvQGVGY iNELLARLTE m.PVRDNTqT NRTLDSSPlT FPLDrSIYAD
A. pediades
                        ALGPVQGVGY VNELLARLTG q.AVRDETQT NRTLDSDPAT FPLNrTFYAD
                   PLGPVQGVGW ANELMARLTR A. FVHDHTCV NNTLDASPAT FPLNATLYAD
P. lycii
A. terreus 9al
A. terreus cbs PLGPvQGVGW aNELIARLTR S.PVHDHTCV NNTLDANPAT FPLNATLYAD A. niger var. awamori PLGPTQGVGY aNELIARLTH S.PVHDDTSS NHTLDSNPAT FPLNSTLYAD
                         PLGPTQGVGY aNELIARLTH S.PVHDDTSS NHTLDSNPAT FPLNSTLYAD
                PLGPTQGVGY aNELIARLTH S.FVHDDTSS NHTLDSSPAT FPLNSTLYAD

PLGPTQGVGY aNELIARLTH S.FVHDDTSS NHTLDSSPAT FPLNSTLYAD

PLGPTQGVGY ANELIARLTH S.FVHDDTSS NHTLDSSPAT FPLNSTLYAD
A. niger T213
A. niger NRRL3135
A. fumigatus ATCC13073 PLGPAQGIGF tNELIARLTR S.PVQDHTST NSTLVSNPAT FPLNATMYVD
A. fumigatus ATCC32722 PLGPAQGIGF 'tNELIARLTR S.PVQDHTST NSTLVSNPAT FPLNATMYVD
A. fumigatus ATCC58128 PLGPAQGIGF tNELIARLTR S.PVQDHTST NSTLVSNPAT FPLNATMYVD
A. fumigatus ATCC26906 PLGPAQGIGF tNELIARLTR S.PVQDHTST NSTLVSNPAT FPLNATMYVD
A. fumigatus ATCC32239 PLGPAQGIGF tNELIARLTN S.PVQDHTST NSTLDSDPAT FPLNATIYVD
                        PLGPAQGIGF tNELIARLTQ S.PVQDNTST NHTLDSNPAT FPLDrkLYAD
E. nidulans
                        PLGPAQGVGF VNELIARMTH S.PVQDYTTV NHTLDSNPAT FPLNATLYAD
T. thermophilus
                        AFGPSRGVGF VNELIARMTG NIPVKDHTTV NHTLDdNPET FPLDAVLYAD
T. lanuginosa
                        PLGPTQGVGF VNELLARLA. GVPVRDgTST NRTLDGDPrT FPLGrPLYAD
M. thermophila
                         PLGPAQGVGF -NELIARLTH S-FVQDHTST NHTLDSNPAT FPLNATLYAD
Consensus Seq. 11
                         FSHDNlmVAV FsAMGLFrqP aPLSTSvpNP wrt....Wr TSSlVPFSGR
P. involutus (phyA1)
                         FSHDNlmVAV FsAMGLFrqS aPLSTSTpDP nrt....Wl TSSvVPFSAR
P. involutus (phyA2)
                         FSHDNqMVAI FSAMGLFNqS aPLdPTTpDP art....Fl vkkiVPFSAR
T. pubescens
                         LSHDNqMIAI FSAMGLFNqS sPLdPSfpNP krt....Wv TSRltPFSAR
A. pediades
                        FSHDNTMVPI FaALGLFNAT a.LdPlkpDe nrl.....Wv DSklVPFSGH
P. lycii
                        FSHDSnLVSI FWALGLYNGT aPLSqTSVES Vs..QTDGYA AAWTVPFAAR
A. terreus 9al
                        FSHDSnLVSI FWALGLYNGT KPLSQTTVEd It..rTDGYA AAWTVPFAAR
A. terreus cbs
A. niger var. awamori FSHDNGIISI LFALGLYNGT KPLSTTTVEN It..QTDGFS SAWTVPFASR
                 FSHDNGIISI LFALGLYNGT KPLSTTTVEN It..QTDGFS SAWTVPFASR
FSHDNGIISI LFALGLYNGT KPLSTTTVEN It..QTDGFS SAWTVPFASR
A. niger T213
A. niger NRRL3135
A. fumigatus ATCC13073 FSHDNSMVSI FFALGLYNGT EPLSTTSVES ak..ElDGYS ASWVVPFGAR
A. fumigatus ATCC32722 FSHDNSMVSI FFALGLYNGT gPLSrTSVES ak..ElDGYS ASWvVPFGAR
A. fumigatus ATCC58128 FSHDNSMVSI FFALGLYNGT EPLSTTSVES ak..ElDGYS ASWVVPFGAR
A. fumigatus ATCC26906 FSHDNSMVSI FFALGLYNGT EPLSTTSVES ak..ElDGYS ASWvVPFGAR
 A. fumigatus ATCC32239 FSHDNGMIPI FFAMGLYNGT EPLSqTSeES tk..ESNGYS ASWAVPFGAR
                        FSHDNSMISI FFAMGLYNGT QPLSmdSVES Iq..EmDGYA ASWTVPFGAR
 E. nidulans
                        FSHDNTMtSI FaALGLYNGT akLSTTeIKS Ie..ETDGYS AAWTVPFGGR
 T. thermophilus
                        FSHDNTMtGI FSAMGLYNGT KPLSTSKIQP ptgaAADGYA ASWTVPFAAR
 T. lanuginosa
                         FSHDNdMMGV LgALGaYDGv pPLdkTArrd ..peElGGYA ASWAVPFAAR
 M. thermophila
                         FSHDNTMVSI FFALGLYNGT KPLSTTSVES I---ETDGYA ASWTVPFAAR
 Consensus Seq. 11
                          mvVErLsC.. fGt...... Tk VRVLVQDQVq PLEfCGgDRn
 p. involutus (phyA1)
                          maVErLsC.. AGt...... Tk VRVLVQDQVq PLEfCGgDQd
 p. involutus (phyA2)
                          mvVErLDC.. GGa......Qs VRLLVNDaVq PLafCGaDts
 T. pubescens
```

```
mvtErLlCQr DGtGsGGpsr imrNgnvQTF VRILVNDaLq PLkfCGgDmd
                     mtVEkLaC.....sgKea VRVLVNDaVq PLEfCGg.vd
A. pediades
P. lycii
                     AYVEMMQCrA ..... ..EK. .EPL VRVLVNDRVM PLHGCPtDKL
                    AYIEMMQCrA ..... .EK...QPL VRVLVNDRVM PLHGCAVDNL
A. terreus 9al
                     1YVEMMQCQA ..... ..EQ...EPL VRVLVNDRVV PLHGCPIDaL
A. terreus cbs
A. niger var. awamori
                     1YVEMMQCQA ..... ..EQ...EPL VRVLVNDRVV PLHGCPIDab
                     1YVEMMQCQA ..... ..EQ...EPL VRVLVNDRVV PLHGCPVDaL
A. niger T213
A. fumigatus ATCC13073 AYfEtMQCKS ...... ..EK...EPL VRaLINDRVV PLHGCDVDKL
A. niger NRRL3135
A. fumigatus ATCC32722 AYfEtMQCKS ...... ..EK...EPL VRaLINDRVV PLHGCDVDKL
A. fumigatus ATCC58128 AYfEtMQCKS ...... ..EK...ESL VRaLINDRVV PLHGCDVDKL
A. fumigatus ATCC26906 AYfEtMQCKS .........EK...EPL VRaLINDRVV PLHGCDVDKL
A. fumigatus ATCC32239 AYFETMQCKS ...... ..EK...EPL VRaLINDRVV PLHGCAVDKL
                      AYFELMQCE. ..... ..KK...EPL VRVLVNDRVV PLHGCAVDKF
                      E. nidulans
                      AYVELLRCET ETSSeEEeEG ..ED...EPF VRVLVNDRVV PLHGCTVDRW
T. thermophilus
                      iYVEkMRCsG GGgGgGGGEG ..rQekdEeM VRVLVNDRVM TLkGCGaDEr
T. lanuginosa
M. thermophila
                      AYVEMMQCEA GG-G-GG-EG --EK---EPL VRVLVNDRVV PLHGCGVDKL
Consensus Seq. 11
                         451
                         GlCtLAKFVE SqTFARSDga GDFEKCFAts a~
 P. involutus (phyA1)
                         GlCaLDKFVE SqAYARSGga GDFEKCLAtt v~
 P. involutus (phyA2)
                         GvCtLDAFVE SqAYARNDge GDFEKCFAt~ ~~
 T. pubescens
                         SICTLEAFVE SQKYAReDgq GDFEKCFD-- --
 A. pediades
                         GvCELsAFVE SqTYAReNgq GDFAKCgfvp se
 P. lycii
                         GRCKrDAFVA GLSFAQAG.. GNWADCF--- --
 A. terreus 9al
                         GRCKrddfve Glsfarag.. GNWAECF--- --
 A. terreus cbs
                         GRCtrDsFVr GLSFARSG.. GDWAECsA-- --
 A. niger var. awamori
                         GRCtrDsFVr GLSFARSG.. GDWAECFA-- --
 A. niger T213
                         GRCtrDsFVr GLSFARSG.. GDWAECFA-- --
 A. niger NRRL3135
                         GRCKLNDFVK GLSWARSG.. GNWGECFS-- --
 A. fumigatus ATCC13073
                         GRCKLNDFVK GLSWARSG.. GNWGECFS-- --
 A. fumigatus ATCC32722
                         GRCKLNDFVK GLSWARSG.. GNWGECFS-- --
 A. fumigatus ATCC58128
                         GRCKLNDFVK GLSWARSG.. GNWGECFS-- --
 A. fumigatus ATCC26906
                         GRCKLKDFVK GLSWARSG.. GNSEQSFS-- --
 A. fumigatus ATCC32239
                         GRCtLDDWVE GLNFARSG.. GNWktCfTl- --
 E. nidulans
                         GRCKrDDFVr GLSFARqG.. GNWEGCYAas e~
 T. thermophilus
                          GRCRrDEWIK GLTFARqG.. GHWDrCF--- --
 T. lanuginosa
                          GmCtLErFIE SMAFARGN.. GKWDlCFA-- --
 M. thermophila
                          GRCKLDDFVE GLSFARSG-- GNWAECFA-- --
 Consensus Seq. 11
```

	M (~~~	·m/~~	****	ייייני	בייים	בידים	כידוכ	ידכנ	TAT	rgc	CAC	CTT	GTT	CGG	T.T.	T CCA	CAI	rcc	3 : 3GT	ACC	20
_	TAC							4				-+-			~-+							60
	A :	L (3 1	?]	2 (2 (2)	G GCT	N ייע מי	S TC1	H	S CTC'	C TTG	D TGA	T CAC	V TGT	D TGA	G ACG	GTG	GT.	Y (Q (C TGT	40
	CGG	_						4				-+-										120
	F TTC	777	ית תרי	יייייי ע	T	CAC	יתיתי	ידיכו	366	TAC	CTA	CTC	TCC	ATP	CI		CI.	110	GCA	CAC	CAL	60 180
121	AAG								<u> </u>			-+-				+			T			100
	man	A	2 MW	m C m	CCD	CAC	V TGT1	רככ	ACD	CGA	CTG	TAC	AGI	TAC	CTT:	TCC	TT	CAA	GT.I	TTG	STCT	80
187												-+-				+					AGA	240
		H ACAC			יא כיז	י ע ער	P CCCI	ממ	ጥጥር	יתיתכ	TGC	CGTC	TA	/GG(CTT.	AC:	CT	GCI	"T.T.C	3W.T.1	E GAA	100 300
241												+ -				+-:					CTT	
					122		T TAC	TCC	بالمالمان	ቦሮ አ 2	Δ CCC	TAI	ΔGT	ACG	CII	L.C.	110	AAC	T BAC	ATT	N CAAC	120 360
301	CG	-+- KTA	GTT	TTT	+ CTT	GCG	ATG	ACG	AA?	AGT	rcc	CAT	rca'	rgc	GAA	MG.	MAC	. 1 1	.10	_	TTG	140
	PO 30	~ ~ ~	mmm/	700	rac	ጥርን	D .CGA		CA	מידירי	CAT	TCG	GTG	AAA	ACC	'AA	ATC	GT	TAA	CTC	G TGGT	140 420
361	AT	-+- GTG	AAA	ccc	ACG	ACT	GCI	'GA	CT	GAG	GTA	AGC	CAC	TTT	TGG	3.T.T.	IAC	.CM	H. I I	O.C.	ncun	160
	-			am a	~ ~ ~	א א כי	Y ATA	מס.	AGG	حبي	TGG	СТА	K GAA	AGA	V TTC	3TT	CCI	TT	CAT	R TAG	AGCT	480
421	 TA	-+- ATT	CAA	 GAT	+ GTC	TTC	TAT	GT	rcc	GAA	ACC	GAT	CTI	TCI	AAC	CAA	.GG	ΓAA	GTA	ATC	TCGA	
					~- ~		nm* =	nma				מממי	ריבועו	1111111	7.T.T.		UUU.	T T T		~~~	A TGCT	
481		_							_ 4 -			+	. – – -			-+-				τ-	ACGA	
	ת ת	CHING.	ccc	מביחיי	CCC	ገልርር	TTT	CTC	AAC	CAC	CACC	'AAC	CT:	CTC	CCA	GT.	LAT	TA	LCG.	I GA:	CALL	200 600
54:	-	_					·									-+-					AGTAA	• • •
	-	1 m C	300	יייי אי	700	كالملك	מהמ	מאמ	מסב	1 CTr	דידיכו	BAC	CAC	GGT.	ACT	"TG	PAC	TG	CLL	E TCG.	AAGAC	220
60	1	+				+			-+-				+			-+				-+-	 CTTC7	

	S	E	L	G	D	D	V	E	A	N	F	T	A	L	F.	Α	بر 	A	 -	R	240
	TCI	'GAJ	\TTI	\GG1	GAC	GAC	CGTI	GAA	\GC1	'AAC	TTC	ACT	GC1	TTC	TTC	:GC1	CCA	AGC.	'AT'	TAGA	~~~
661		-+	- -		-+-							+			-+-						720
	AGA	CT	CAA7	rcc <i>i</i>	CTC	CTC	GCAP	CTI	CGI	TTC	JAAG	TGA	CGA	AAC	CAAC	CGF	\GG:	rcg	ATA	ATCT	
	A	R	L	E	A	D	L	P	G	V	T	L	T	D	E	D	V	_V	Y	L	260
	מכיי	ואמי	ነጥጥ <u>ር</u>	GGAJ	AGC"	rga(CTTC	GCC#	AGG:	rgT?	raci	TTC	ACT	GAC	CGA	AGAC	GT:	rgt'	TTAC	CTTG	
721					+-							+			+-				+		780
,	CGZ	ATC'	raa	CCT:	rcgi	ACTO	JAAC	CGG:	rcci	ACA	ATGA)AA	TGF	CTC	CT:	CTC	3CA	ACA:	AATO	BAAC	
																					-
	М	D	M	C	P	F	D	T	v	A	R	T	S			T			S		280
	70.000	~~~	ጣ አ ጥ ሰ	- ಇಗಿದ್ದ	דככ	ידים	CGA	ים בר	TGT	cgc'	TAG	AACI	rrc:	GAO	CGC'	rac:	rga.	TTA	GTC:	rcca	
781	<u> </u>				+				+			-+			+				+		840
, , ,	TA	CCT	GTA	CAC	AGG'	TAA	GCT	GTG	ACA	GCG:	ATC?	rtg <i>i</i>	\AGI	CT	GCG	ATG	ACT'	TAA	CAG	AGGT	
	F	C	A	L	F	Т	Н	D	E	W	I	Q	Y	D	Y	L	Q	S	L	G	3 00
	фф	ста	TGC	TTT	GTT	CAC'	TCA	CGA	CGA.	ATG	GAT	CA	ATA	CGA	CTA	CTT	GCA.	AAG	CLLL	GGGT.	
847					+				+			-+-			+				+		900
011	AA	GAC	ACG.	AAA	CAA	GTG.	AGT	GCT	GCT	TAC	CTA	GT.	TAT	3CT	GAT	GAA	CGT	TTC	GAA	CCCA	
	ĸ	Y	Y	G	Y	G	A	G	N	P	L	G	P	A	Q	G	V	G	F	A	320
	ממ	at D	ርጥል	CGG	TTA	CGG	TGC	TGG	TAA	CCC	ATT	GGG'	TCC.	AGC	TCA	AGG	TGT	TGG	T.T.T.	CGCT	
901									+			-+-			+				+		960
	TT	CAT	GAT	GCC	AAT	GCC	ACG	ACC	TTA	GGG	TAA	CCC.	AGG'	TCG	AGT	TCC	ACA	ACC	AAA	GCGA	
	N	E	L	I	A	R	L	T	H	S	P	V	Q	D	H	T	S	T	N		340
	AA	CGA	ATT	GAT	TGC	TAG	ATT	GAC	TCA	CTC	TCC	AGT	TCA	AGA	CCA	CAC	TTC	TAC	TAA.	CCAC	
961					4				+			-+-			+				· — —		1020
	ТT	GCI	TAA	CTA	ACG	ATC	'T'AA	CTG	AGI	GAC	AGG	TCA	AGT	TCI	'GGT	GTG	AAC	ATC	ATT	GGTG	
																					360
	T	L	D	S	N	Þ	A	T	F	P	L	N	A	T	ь	Y.	A	עם	F.	S	360
	AC	TTT	GGA	CTC	TAA	CCC	:AGC	TAC	TTT	CCC	TTA:	'GAA	.CGC	TAC	TTT	GTA	CGC	TGF	rc.L.T	CTCT	1000
1021												-+-			+						1000
	TO	AA	CCI	GAG	ATI	GGG	TCC	SATO	LAAE	\GG(STAA	CTI	'GCG	ATC	AAA	CAT	GCC	AC.	GAF	GAGA	
													•								380
	H	D	N	T	M	I	S	I	F	F	A	L	G	L	Y	N	ن مصح	T	K		380
	CF	ACGI	CAA	ACAC	'TAT	CAD'	TAT	CTA	rtt'	rct'	rcgc	TTT	'GGG	TT	rgta	CAP	(CGC	3.T.Y.	CAF	AGCCA	7140
1081		4			4	L			-+-			+-									TIFO
	GT	rgc'	rgT?	rgte	AT	ACT#	ATA	GAT!	AAA	AGA	AGCC	IAAF	CCC	'AA'	ACA'	rgt	rgc	CAT	3GT.	rcggt	
																					400
	L	S	T	T	S	V	E	S	I	E	E	T	D	G	Y	S	A	S	w ~~~	T TCDCT	400
	-		7077	7077	amm/	ייייייי	בסתייו	አ አ ጥ	ומידים	ביתית	מאמ	1 A A L	TGP	\CG(31'17	IC.I.(TIG	-11	-11	3GWC I	
1141	1					L ·			- 4 -			+-				+			-+-		1200
	AZ	ACA	CAT	TAE	AAE	3AC	AAC'	TTA	GAT.	AAC	TTC	rtt	GAC!	rgc	CAA!	rga	GAC	GAA	GAA	CCTĠA	
	v	P	F	A	A	R	A	Y	v	E	M	M	Q	C	Q	A	E	_ K	E	P אארכא	420
			~ > ~	~~~	ama (OMB /	~~~		700	ጥጥር	ימממ	ימבאי	rgcz	TA	GTC.	AAG	CTG.	MMM	MGG	wece	
120												+				+					
	C	AAG	GTA.	AGC	3AC	GAT	CTC	GAA	TGC	AAC	TTT.	ACT	ACG'	ГTА	CAG	TTC	GAC	TTT	TCC	TTGGT	
	L	v	R	v	L	v	N	ם	R	: V	v	P	L	H	G	С	A	. v	D	K NCABG	540
	-		~~~	~~~	THE PERSON NAMED IN	$m \sim c^{-\epsilon}$	V ethan	700	מים תי	CAC	שיייים	ጥጥር	ሮልጥ	TGC	ACG	GIT	GIG	c_{1}	1770	VCVVC	•
126	-											+				+					
	A	ACC	AAT	CTC	AAA	ACC	AAI	TGC	TGT	CTC	:AAC	AAG	GTA	ACG	TGC	CAA	CAC	GAC	!AAC	TGTTC	:
	L	G	R	C	ĸ	R	I) I	, ,	E	G	L	S	F	A	R	2 5	; G	G	460

1201									TTCGTTGAA			
1321									BAAGCAACTT			
	N	W	A	E	С	F	A	*	467			
								TTA	1410			

TTGACCCGACTTACAAAGCGAATT

1	M G	;	V	F.	v	V :	L	L	S	I	A	T	L	F GTT	G CGG	S	T CAC	S TAT	CC0	GT	r acc	20
	ATGC TACC					_	1			4				-+-			7					60
	A I		G	P P	R	G 'GGT	N 'AAC	S TCI	H	s TC1	C TG:	D rga	T CAC	V TGT	D TG#	G ACG(G STG0	Y TTE	AC(Q CAA	C TGT	40
	CGG			. .			-4-							-+-								120
	F TTC	P CCA	ממט	יידע	וייירים	rcac	TTC	TGC	3GG	rac:	ATA	CTC	TCC	LTA	CT	LC.L	C.L.T.	Lec	CT	ع بری	E GAA	60 180
				•							+			-+-				+			T	100
	S .	~~"	ייתו איי	ייייייי	TOO	222	<u> ገርጥ</u>	rcc:	AAA	GGG'	${ t TTG}$	TAG	AGT	"I'AC	T.T.	TCG	TIC	WWG	3 T T	110	11	80
181														· - + -				+			AGA	240
				mac	mn c	Y ATA		אמ מ	ጥጥር	ጥጥር	TGC	GTC	TAT	いいしょ	ـ تات	WC T	CIG					100
241				_			1				+			+:				-			ACTT	300
					~~ A	A CGC	ጥአሮ	TCC	لململما	ממיזי	(GGC	TA	AGT	ACG	$c_{1,1}$	101	. IGA	M.G.	~~·			120
301									_ _					+				7			+ GTTG	360
	Y TAG			~~~	-ma	D CTGA	CON		ו מבחת	יידרי	ידער	rcg	GTG	AAC	AAC	AA:	I V	STT	AA	CTC	TGGT	140 420
361											- 4 -			+				•			ACCA	
					2026	R SAAC	ידו אור	י מים ו	אככו	بالماليات	TGG	CTA	GAA	AGP	7.T.T.C	21.1.	CCA	776		17.0	122002	160 480
421								L			-+-										TCGA	
				~~~	202	V GAG'	וגידור	rrc	سلسل	<del>ረ</del> ምር	CTC	ΔΔ.	\AG'I	TC	7.T.T.	GMM	r DD					180
481				_											+						+ GACGA	
						G CAG			700	יראר	זים מי	ואאי	3C."I"	CIT	ししみ	CI	Lar.		~~.			200 r
54	_								<b>-</b>		4 .				+			+			ATAA	•
	P	E	: G	3 7	A (	y	N	ı N	1 7	נ ז	. ]	באר.	H (	G GGT	L TTC	C TG:	T TAC'	A TGC	F TT:	E TCG	E AAGA	220 A
60	_								<b></b>						+						TTCT	

	S	E	L	G	D	D	V	E	A	N	F	T	A	V	F	A raco	P	) C.C.	_	R 'DGD	240
			TTG	GGT	GAC	GAC	GTI	'GA	\GC'	CAAC	CTTC	ACT	.GC1	. <del></del> -			+-			AGA	720
661	AGA	CTT.	'AAC	+	CTG	CTC	CAZ	CTI	rcgi	ATT	GAAC	TGP	CGF	CAZ	AA/	GCG?	AGG	rgg	TAP	TCT	
	A	R	L	E	A	Н	L	P	G	V	N	L	T	D rgag	E	_	V EGT:		-	L TTG	260
721				. 4			+-				+			-+			+			+	780
	CGA	TCT	OAA1	CTI	CGA	GTO	)AAE	CGG'	rcc.	ACA	ATT(	)AAE	CTG	ACTO	GCT:	rct(	3CA	ACA	ATTO	BAAC	-
	M	D	M	C	P	F	D	T	v	A	R	T	S	D	A	T	Q TCA	L	S	P.	280
781		. – – .		. <b>.</b>			+				+			-+			+			ACCT	840
	TAC	CT	STAC	CACA	\GG1	CAAC	3CT(	GTG.	ACA	ACG.	ATC'	L.L.C.	AAGA	ACT	366	MIG.	MGI	I TARN		AGGT	
	F TTO	C	D TGA(	L TTC	F STT(	T CAC	H TCA	D CGA	E CGA	W ATG	I GAT'	Q TCA	Y ATA	D CGA	Y CTA	L CTT	Q GCA	S ATC'	L TTT	G GGT	300
841				-+			+		<u>-</u>		+			-+-			+			+	900
	AA	EAC	ACTO	)AAE	CAAC	GTG	AGT(	GCT	GCT	TAC	CTA	AGT"	rarc	3C.T.	GA'I'	GAA	CGI	I AUJ	H4444	CCCA	
	K	Y	Y CTA	G CGG'	Y TTA	G CGG'	A TGC	G TGG	N TAA	P .CCC	L ATT	G GGG'	P TCC	A AGC	Q [.] TCA	G AGG	V TGT	G TGG	F TTT	V CGTT	320
901					~ .		+				+			-+-			+			+	960
	TT	CAT	GAT	GCC	AAT(	GCC.	ACG.	ACC	ATI	'GGG	TAA	CCC.	AGG	TCG.	AGT	TCC	ACA	ACC	MAN	GCAA	
	N	E	L	I Candri	A	R TAG	L ልጥጥ	T	H TC2	S	P TCC	V AGT	Q TCA	D AGA	H CCA	T CAC	S TTC			H CCAC	340
961				-+-			+				+			-+-			+				1020
	TT	GCT	TAA	CTA	ACG.	ATC	AAT	.CTG	AGT	GAG	AGG	TCA	AGT	TCT	GGT	GTG	AAG	ATG	ATI	GGTG	
	T	L	ם	s	N	P	A	T	F	P	L	N GND	A CGC	T TAC	L TTT	Y GTA	A .CGC	_	F .CTT	S	360
1021							+				+			-+-			+			+	1080
	TG	AAA	.CCT	GAG	ATT	GGG	TCG	ATC	IAA	\GGG	TAA	CTT	GCG	ATG	AAA	CAI	GCG	AC'1	GAA	GAGA	
	H	D	N	T	M	٧	s	I	F	F		L							K ממחיי		380
1081							4				- +			-+-			+			GCCA	1140
1001	GT	GCI	'GTI	GTG	ATA	CCA	AAC	ATA	LAA	AGAJ	AGCG	AAA	CCC	'AAA	CAI	GTI	GCC	CATO	TA	CGGT	
	<b>.</b>	_	m	m		17	-	c	т	E	E	т	D	G	Y	S	A	s	W	T	400
•	ma		יתים כ	ישא ר	יתיתיר	יייים	ל ביורייו	ነ አጥር	י עיטיב	TTG?	AAGI	<b>VAA</b>	TGA	CGC	TTT	CTC	TIGO	LLLC	TIT	3GAC I	
1141							4				-+			+ -						CTGA	1200
																	-				
	ν	p	F	A	A	R	A	Y	v	E	М	M	Q	C	E	A	E	X	E	P	420
1001	GI	TCC	AT!	rcgc	TGC	CTAC	GAG	CTT	ACG'	TTG:	AAA.	rga:	rgcz	\AT(	3TG/	4AG(		144. 		AACCA +	1260
1201	. CP	AGC	TAI	AGCC	ACC	TAE	CTC	'AAE	rgc	AAC'	TTT	ACT	ACG7	CTA(	CAC	rtc	JAC:	r <b>t</b> t'	TCC'	TTGGT	
	L	v	R	v	L	v	N	D	R	v	v	P	Ļ	H	G	С	G	v	D	K	440
	thu:	can	ו עידיין	באמי	יישיטים	ימכי	מידיד	DO A	ACA	GAG	TTG'	TTC	CAT	rgc	ACG(	$\mathtt{GTTC}$	GTG	SIG	TTG.	ACAAC	; - 1320
1261	A.F	CC	LAT	CTC	AAA	ACC	TAA	TGC	TGT	CTC	AAC	AAG	GTA	ACG'	rgc	CAA	CAC	CAC	AAC	TGTT	:
	L	G	R	С	к	R	D	D	F	v	E	G	L	s	F	A	R	S	G	G	460

WO 99/48380 PCT/DK99/00154

1321	AACCCATCTACATTCTCTCTGCTGAAGCAACTTCCAAACAGAAAGCGATCTAGACCACCA											
								* TTAA	467			
1381				-+-			+		1404			

	M	G	v	F	v	v	L	L	s	I,	A	T	L	F	G	. S	T	S	G ZGG(	T TACT	20
1											+			-+-			+			CACT + ETGA	60
	TAC	:CCC	CAF	AAC	3CAC	i CAA	AGA.	LAA.	LAGA	41W	GCG	C 1 C.	1022								
	CCC	ירייר	2000	יככנ	CG	rggz	'AAA	TCA	CTC	CAA	GTC	CTG	CGA'	TAC	GGT.	AGA	CCT.	AGG	GTA	Q CCAG	-
61				- +			+				+			-+-			+			GTC	120
	c c	S	P	A	T	S	H	L	W	G	T CAC	Y GTA	S CTC	P GCC	Υ ATa	F CTT	S TTC	L GCT	E CGA	D GGAC	60
121							+				+			-+-			+			+ CCTG	180
	E	т,	s	v	s	s	ĸ	L	P	ĸ	D	С	R	I	T	L	v	Q	v	L	80
181	GA	GCT(	GTC	CGT	GTC	GAG'	TAA	GCT	TCC	CAA	.GGA 	TTG.	CCG	GAT -+-	CAC	CTT	+ GG.T.	ACA		+	
	CT	CGA	CAG	GCA	CAG	CTC.	ATT	CGA	AGG	GTI	'CCI	AAC	:GGC	CTA	.GTG	GAA	CCA	TGT.	CCA	CGAT	
	TO	200	רראי	TGG	AGC	GCG	GTA	CCC	AAC	CAC	CTC	CAA	GAG	CAA	AAA	GTA	TAA	GAA	GC.T.	I TaTt	
241	AG	 CGC	GGT	-+- ACC	TCG	CGC	CAT	GGG	TTG	GTC	GAG	GTI	CTC	GTI	TTT	CAT	'ATI	CTT	CGA	+ AtAa	
	3.0	~~~	$\sigma$	CCN	aac	ממס	TGC	יר א ר	CGA	רידים	CAZ	\GGC	CAP	\GTa	LCGC	:CT.T	TIT	GAH	CAL	Y GTAC	
301											. +			+-						+ CATG	360
	NT	v	ηı	. т.	æ	<b>7</b> A.	מ	מ	Τ.	T	P	F	G	E	Q	Q	L	v	N	s	140
361	AA	CTA	TAC	TCT	'GGG	TGC	:GGA	TGA	CCI	CAC	TC(	CCT:	rtgo 	3GG <i>F</i> +-	4GC/					+	420
																				GAGC	160
	CC	ר מים	מ מיזי	CTT	מידטי	CCA	GAG	GTA	CAZ	\GG(	CTC'	rgg	CGC	<b>JCA</b> (	FTG:	I'GG'	ruc	1011	TWI	R TCGC	
.421	CC	GTA	GTI	CAA	GAI	GGI	CTC	CAT	GT	CCC	GAG	ACC	GCG	CGT	CAC	ACC!	ACG(	GCA <i>I</i>	\ATI	AAGCG	
			33.00	10mc	2007	CCC	2007	ם עודה	יידיכו	ישישיי	רככ	にかば	AGA:	AGT"	TCA'	TCG	AGGG	. لا فاد	r	Q AGCAG	
483							4	L			-+-			+				+		rcgro	240
	25.	ĸ	τ.	A	Ð	P	G	A	т	N	R	. A	. A	P	A	I	s	v	I	ı	200
54:	1							<b>.</b>			-+-			+				+		TTATT + AATAA	. 600
																				AATAA A	
<i>-</i>	C	rac:	מממ	3061	227	ملتات	TCA	ACA	ATA	CGC	TGG	ACC	:ACG	GTG	TGT	GCA	CGA	$\mathbf{AGT}$	TTG.	AGGC	3
60	. G(	GCC'	TCT	CGC:	rcro	GCA	AGT'	TGT'	TAT	GC©	ACC	TGC	TGC	CAC	ACA	CGT	GCT	TCA	AAC	TCCG	2

				G GGA			-mm		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	יתתי	ידידידיד	: זביי	166	ヒーコ	. •		-				R CGA	720	
				+														•				120	
	A	R	L	E	K	Н	L	P	G	V car	T GAC	L	T 'GAC	D AG	E ACG/	D AGG	V ACG	V TTC	TC	S :	L CTA	260	
721													CTC	TC:	rgc'	rcc	TGC	AAC	'AG	TCA	GAT		-
781	AT	3 <b>G</b> A	CAT		rcC	GTT'	TGA:	rac	GGT	AGC	:GCG	CAC	CAC	3CG		CAA	GTC	:AG(	CTG	TCA 	P .CCG +	28¢	_
781	TA	CCT	GTA	CAC	AgG	CAA	ACT!	ATG	CCA	TCC w	CGC x	GT(	GT( Y	CGC D	TGC Y	. I	i (	) }1C	sac S	L	G		0 .
841	TT	CTG	TCA	ACT	CTT	CAC	TCA	CAA	TGF	GT	3GA/	AGA.	AGT.	AC9 +	ACI			-+-			GGC + CCCG	90	0
	ĸ	Y	Y	G	Y	G	A	G	N	P	L	G TGG	q GAC	A CGG	L C	) (	g GGG	I ATA	.GG(	F 3TT	T CACC		
901							GCG	TCC	GT	rgg	-+- GAG	ACC	CTG	GCC	GAC	TC	CCC	TAT	CC	CAA	GTGG		340
961	A	ACG.	AGC'		rtgo	ccc	GTI		CgC	GTT	CGC	CAC	TGC	AGC	BAC(	CAC 	ACC	AGC	:AC		CTCG	. 10	
,,,,	T'	rgc	TCG.	ACT	AAC	GGG	CCA	ACTO	GCG	CAA	GCG	GTC	ALC	3 I C	-10	J					,		360
102	A	CTC	TAG		CCA	ACC	CGG	CCA	CCI	TCC	CG1	TG	AAC	3C1/	4			+			TTC		080
				_				_	-	, ,	. z	٠ .	r. (	G	L	Y	N	G	T	E	P		380
108	C	ACC	ACA	ACA	GCA	TGG	TTT	CCA	TCI	TC:											ACC		140
	I	؛ ر	3 I	<b>२</b> 7	: s	s v	7 E	. S	S 2	A :	K :	E	L	D	G rece	Y Ta'	S TTC	A TGC	S AT	W CCT	V GGGT	G	
114	1 -	AAC	AGG	GCC	rgg <i>I</i>	AGC	CACC	+ TT	rcg	cgg	+ TTC	CTI	AAC	CTA	CCC	CAT	AAG	ACC	TA	GGA	CCCA	.C	
	(	GTG	CCT	TTC	3GC(	GCG	CGAC	3CC	rac	TTC	GAG.	ACC	ATC		-+-				+		AGCC	+	
120		CAC	GGA	AAG	CCG	CGC	GCT	CGG.	ATG	AAG	CTU	TGC	TAC	_G L	1320	911		,,,					
12		CTI	GTI	CGC	GCT	TTG	ATT.	AAT	GAC	CGC:	sGT".	r			-+-				+	. <b></b> -	KEACAI	-+	1320
		GAJ	CAP	'GCG	CGA	AAC	TAA	TTA	CTC.	GCC	CA	ACA	دوو	IGM	CGI	nc.							460

							TTTGTCAA						• ^
1321							AAACAGTI					136	_
		 _			S TAG	* TTGA	467						
1381	-:-	 	-+-	 	+		1404						

	CP-1 Eco TATATG	3 3 TTC	M G <u>atg</u> gg	CCTC	rTCGT	CGT	CTA	CTGI	CCAI	TGC	CACC	TIG	TTC	267.1	CLA	
1	ATATAC						-+					+				60
		T	A L	G :	P R	G TTGG	N TAAT	S F	I S	C CTTG'	D TGAC	T CACT	V :	D (	G GGTG	120
61	GTAGGC C	CATGG P-2	CGGAA	CCCA	GGAG(	CACC.	ATTA	AGA	TGA	SAAC.	ACT	3 I GM	,ÇAA	C10.	-cac	
121	GTTACC	AATGI		AGAA	ATTT	CTCA	CTTG -+-	TGG	GTC:	AATA +	CTC:	rcca +	TAC		+	180
121	CAATGO	TTAC	AAAGGG	TCTT	<b>KAAT</b>	GAGT	GAAC	'ACC	CCAG'	TTAT	GAG	AGGT	AIG	AAG.	MGAA	
181	TGGAA	BACGAI		CTATI	TCTC	CAGA	CGT	CCA	GACG	ACTG +	TAG	AGTI +	ACI	TTC	<del>-</del>	240
	ACCTT	CTGCT'	ragac(	CP-4	lagag <u>L.7</u>	GTC1	GCA	AGGT	CTGC	TGAC	!ATC	TCA	ATGA	LAAG	CAAG	
241	AAGTT'	t-	TAGAC		A F	Y GATA	ACCC:	aact 	gacT	CTA?	⊾Ggg 	taa	3aa9 +		+	300
241	TTCAA	aacag.	ATCTG'	TGCC	ACGAI	CTA:	rggg'	TTGA	.ctgA	GAT	rCcc	aTT	Ctto	EATG	AGAC	į
201	CTTTG		AGCTA		AAAG	AACG	CTAC	TGCI		AGG(	STAA 	GTA	CGC: +		:TTGA +	360
301	GAAAC	TAACT	TCGAT	AAGT'	TTTC:	TTGC	GATG P-6	ACGA	LAAGI	rtcc	CATI	CAT	GCG:	DAAA	;aact	•
	T AGACT	מ מיז מידי	Y T ACATO.	CTTT	GGGT	GCTG	D ACGA	L CTT	T I	CCAT	TCG	TGA	h-wa	CCA	AATGO	
361	TCTGA	ATGTT	GATGT	GAAA	CCCA	CGAC	TGCI	'GAA	CTGA	GGTA	AGC	CACT	TTT	GG1.	IIAC	
43:	TTAAC	TCTGG	I K	AGTT	CTAC	agaa 	GAT?	CAA	 3GCT'	TTGG -+	CTA		+			+ 480
42.	OTTAA	BAGACO	TAATAS	TCAA	GATG	TCTI	CTAT	rgtt:	CCGA <u>CP-</u>	8.7 CF	-9	CITI			AUU I	•
4.0	TCAT	PAGAGO	s (	GTTC	Ttct	AGAC	3TTA'	rtgc 	TTCT	GCTG	;aaa 	AGT.	-+			- + 540
40	AGTA	ATCTC	GAAGA(	CAAC	SAaga	TCT	CAAT.	AACG	AAGA	CGAC	.111	ICA	, G _ *	2101		
E A	TCCA	ATCTG	K 1 CTAAG	TTGG	CTGA	CCA	ggtt +	CTCA	ACCA	CAC	CAAG		-+-			+ 60
J-1	AGGT	TAGAC	GATTC.	AACC	GACT	GGT	CCAA	GAG'	TGGT	rgtg	GTTC	GAA	GAG	GTCF	ATA	C

WO 99/48380

## 29/32

	<u>CP-10.7</u>																				
CP-11.7 V I I <u>S</u> E A S S Y N N T L D P G T C T												7	•								
	v	I	I	s	E	A	s	s	Y	N	N	T	L	D	P	G	T	C	T	A	
	ACGT	TAT	TAT	Tto	tGA	cgc	tTC	Ttc	tTA	CAA	CAA	CAC	TTT	GGA	CCC	aGG	TAC	TTG	TAC	TG	
601			+				+			-+-			+				+			-+	660
	TGCA	ATA	ATA	Aag	aCT	'gcg	aAG	Gag	aAT	GTT	GTT	GTG	AAA	CCI	'Ggg	tCC	ATG	AAC	ATG	AC	

T: .. 40D

61	CTI	TCG	AA	GAC	TC:	rgal	ATT	Ggc	tGA	Cac	EG1	167		+				L TTT( +			rc -+ '	720	
61	GAJ	AAGO	TT	CT	BAG	ACT	ľAA	ceg	acı	Gug	aur								CP-	12.	<u>7</u> .		
	CA	GCT:	AT I	'AG	AGC	TAG	ATT	'GGA	AGC	TGP	rc.r.ı	. با	_AGC		h			. 4		E tga	-+	780	-
721	GT	CGA	TA	+ \TC	TCG	ATC	TAA	CCI	TCG	ACT	rgai	ACG(	3TC	CACI	AAT	iaae	CTG	BACI	Gtg	ract	tC.		
	TI	act	TA	CTI	'GA'	rggz	CA?	rgro	Tt	CET	T.C.G.				+			-+-		T CTAC	+	840	
781	A.	Ltga	LAT	GAI	CT	ACC:	rgiz	ACA	LAd	yan	AGC.												
		ATT(	FTC	TC	CAT'	rcr(	GTG	CTT	TGT	TCA	CTC	ACG	ACG		+			-+-		L ACT'	+	900	)
841	T'	AAT	CAC 2	AG P-	GTA 14.	aga Z	CAC	GAA	ACA	AGT	GAG	1160											
	A	ATC	TT:	гGа	K agA	AGT	Y ACI	ACG	GTC	acc	3GTG	SC-T-C			- +			+-		:	V GTG	960	0
90:	1 - T	TAG	AA	ACt	+ tcI	TCA	TGA	TGC	CAS	gtg(	CAC	CGA(	CCA'	rtg(	<b>GT</b>	AAC	CAC	GTt Q I	gac	TTC	CAC	2	
0.0		TGG	TT	TC	CT	AAC	AA:	rtg	TTA	GCT.	R : AGA:	TTG.	ACT.	AGA'	TCT	CCA	GTT 	CAA(	GAC	CAC!	ACTI +	r + 10	20
96	1 -					rtg	CTT: C:	AAC' P-1	FAA 6 	CGA	TCT.	AAC	TGA	TCT	AGA	GGI	<u> </u>						
	ı					- ~~		D .	S	N	P	(30.71	ALL					-			Y Z	A G + 10	080
102	21	 GAT	GAT	 CTG	-+- GTG	 TGA	AAC	+ CTG	AGA	TTG	GGI	CG?	TG	AAC	GGT	CAAC	TTC	CGA	TGA	AAC	ATG	C	
•		CTG	AC'	TTC	TCI	CAC	:GAC	'AAC	ggt	cati	CAT.	LIC.			+				+		N	+ 1	140
10	81	GAC	TG:	AAC	AGA	KGTC	CTC	31.10	3CC	ıLa.	CP-	18.	7	0 7									
		GT	ACT	GC'	rcci	TTA	GTC'	TAC'	TAC	TTC	TGT	E TGA	S ATC	I TAT	E TGA				+		S CTC	-+ 1	200
11	.41	CA'	rg?	CG.	AGG	TAA	CAG	AIG	AIG	MAG	AC.												
		ct	gcl	TG	GAC	TGT	TCC	'ATT	Cgc	ttc	: tag	AGC.	71.13						-+		A	-+ :	126
13	201	ga	.cg	aAC	+ CTG	ACA	AGG	<b>AAT</b>	.Gcc	jaaç	gaTC	CTC	GAA:		\AC1 ?-2(	_	ACT	ACGT	)AT1	CAGI	rcg	iAC	

						R															
	AAAA																				
1261	TTT	CCI	+ TGG	TAA	 CCA	ATC	+	 AAA	.cca	-+- TTA	GCI	GTC	TCA	ACA	agg	TAA	.CGT	GCC	AAC	AC	1320

												F								
1321	CTGT		+				+			-+-			+		 	+			-+	1380
	GACA	ACT	GTT	CÁA	.ccc	ATC	TAC	ATT	CTC	TCI	GCI	'GAA	GCA	ACT	AAA P-2		AAA	.GCG	AT	
	GATO	TGG:	TGG	TAA	CTG	GGC	TGA	ATG	TTT	'CGC	TT		ATT	CAT						
1381	CTAG															26				-

#### INTERNATIONAL SEARCH REPORT

International application No. PCT/DK 99/00154

#### A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A23K 1/165, A01H 5/00, C12N 9/16
According to International Patent Classification (IPC) or to both national classification and IPC

#### **B. FIELDS SEARCHED**

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A23K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

c. Docu	MENTS CONSIDERED TO BE RELEVANT	
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Х	WO 9716981 A1 (GIST-BROCADES B.V.), 15 May 1997 (15.05.97), See Example 2, page 10; and the claims	1-4
	<del>"-</del>	
х	EP 0619369 A1 (AVEVE N.V.), 12 October 1994 (12.10.94), See page 5, lines 11-15; page 7, lines 1-3; and claims 20-21	1-14
х	EP 0682876 A1 (SOUFFLET ALLMENTAIRE), 22 November 1995 (22.11.95), See page 3, lines 21-25 and claim 9	1-8
		·
х	WO 9114782 A1 (GIST-BROCADES N.V.), 3 October 1991 (03.10.91), See page 2, lines 25-27	9-14
		·
<u> </u>		

	(03.10.91), See page 2, line	s 25-	27
			·
X	Further documents are listed in the continuation of Box	C.	X See patent family annex.
* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" "L"	erlier document but published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other	"X"	document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"O"	special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than	"Y"	document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person scilled in the art
- P	the priority date claimed	"&"	document member of the same patent family
Dat	e of the actual completion of the international search	Date	of mailing of the international search report
6	July 1999		<b>1 3</b> -07- ₁₉₉₉
	ne and mailing address of the ISA/	Autho	rized officer

Carolina Palmcrantz/Els

#### INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 99/00154

		FC1/DK 33/C	,013 <del>-1</del>
C (Continu	ation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Gtation of document, with indication, where appropriate, of the relevant	ant passages	Relevant to claim No.
A	WO 9735016 A1 (NOVO NORDISK BIOTECH, INC.), 25 Sept 1997 (25.09.97), See page 41, line and claim 40	s 7-8;	1-14
A	Dialog Information Services, File 5, Biosis, Dialog accession no. 10902627, Biosis acce no. 199799523772, Jiang Junping: "Thermost phytase from Aspergillus sp.", Weishengwu Xuebao 36 (6): p476-478 1996	ssion able	1-14
	<del></del>		
		, i	!
			·
	·		
	•		-

#### INTERNATIONAL SEARCH REPORT

Information on patent family members

01/06/99

International application No.
PCT/DK 99/00154

	document	.	Publication		Patent family		Publication	
cired in s	earch repor		date		member(s)	•	date	
WO 9	716981	A1	15/05/97	AU	7567696	Δ	29/05/97	
.,			-5, 40, 51	BR	9606683			
							09/06/98	
				CA	2209010		15/05/97	
				CN	1168084		17/12/97	
				CZ		A	18/03/98	
				EP	0804087	Α	05/11/97	
			•	HU	9801179	A	28/08/98	
				IL	121208	D	00/00/00	
				JP	10512456	T	02/12/98	
				PL	321186	A	24/11/97	
		•		SK	90797		06/05/98	
			18/18/04					
EP 06	519369	A1	12/10/94	CA	2120265		06/10/94	
		*		FI		A	06/10/94	
				JP	6319539	À	22/11/94	
				NO		A	06/10/94	
				US	5443979		22/08/95	
				US	5554399	A	10/09/96	
EP 06	 582876	A1	22/11/95	AT	172 <b>07</b> 8	т	15/10/98	
_, 50	,	***	mm/ A A / 3/J	DE	69505299			
			•	FR	2719978		00/00/00 24/11/95	
						ハ, ii 		
WO 91	14782	A1	03/10/91	UA	632941	В	14/01/93	
				AU	649447	В	26/05/94	
				AU		A	21/10/91	
				AU		Ā	21/10/91	
				CA	2054762		24/09/91	
				CA		Ä	24/09/91	
•				EP		Ä	02/10/91	
				EP	0449376		02/10/91	
				FI	915477		00/00/00	
				FΪ		D	00/00/00	
				HŪ	215164		28/10/98	
				HU	215260			
				IL	97645		30/11/98 18/03/97	
				11_	3/043	<b>A</b>	10/03/3/	
				JP	6501838	T	03/03/94	
				JP .	6501838 6502296	T T	03/03/94 17/03/94	
				JP JP PT	6501838 6502296 97110	T T A,B	03/03/94 17/03/94 29/11/91	
ı				JP JP PT PT	6501838 6502296 97110 97111	T T A,B A	03/03/94 17/03/94 29/11/91 31/12/91	
,				JP JP PT PT US	6501838 6502296 97110 97111 5543576	T T A,B A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96	
				JP JP PT PT US US	6501838 6502296 97110 97111 5543576 5714474	T T A,B A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96 03/02/98	
				JP JP PT PT US US	6501838 6502296 97110 97111 5543576 5714474 9114772	T T A,B A A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96 03/02/98 03/10/91	
				JP JP PT PT US US WO US	6501838 6502296 97110 97111 5543576 5714474	T T A,B A A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96 03/02/98 03/10/91 14/01/97	
				JP JP PT PT US US	6501838 6502296 97110 97111 5543576 5714474 9114772	T T A,B A A A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96 03/02/98 03/10/91	
	 35016		25/09/07	JP JP PT PT US US WO US	6501838 6502296 97110 97111 5543576 5714474 9114772 5593963 5770413	T T A,B A A A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96 03/02/98 03/10/91 14/01/97 23/06/98	
WO 97	35016	A1	 25/09/97	JP JP PT PT US US WO US US	6501838 6502296 97110 97111 5543576 5714474 9114772 5593963 5770413	T T A,B A A A A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96 03/02/98 03/10/91 14/01/97 23/06/98	
WO 97	35016	A1	 25/09/97	JP JP PT US US WO US US	6501838 6502296 97110 97111 5543576 5714474 9114772 5593963 5770413 	T T A,B A A A A A A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96 03/02/98 03/10/91 14/01/97 23/06/98	
WO 97	 35016	A1	 25/09/97	JP JP PT US US WO US US AU AU	6501838 6502296 97110 97111 5543576 5714474 9114772 5593963 5770413 2077197 2539197 2248980	T T A,B A A A A A A A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96 03/02/98 03/10/91 14/01/97 23/06/98 	
WO 97	35016	A1		JP JP PT US US WO US US AU AU CA EP	6501838 6502296 97110 97111 5543576 5714474 9114772 5593963 5770413 	T T A,B A A A A A A A A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96 03/02/98 03/10/91 14/01/97 23/06/98 	
WO 97	35016	A1		JP JP PT US US WO US US AU AU	6501838 6502296 97110 97111 5543576 5714474 9114772 5593963 5770413 2077197 2539197 2248980	T T A,B A A A A A A A A A	03/03/94 17/03/94 29/11/91 31/12/91 06/08/96 03/02/98 03/10/91 14/01/97 23/06/98 	

	·		
	·		
			· .
·		· ,	
·			
	·		
			٠